

Protocol

Achieving Consensus in the Development of an Online Intervention Designed to Effectively Support Midwives in Work-Related Psychological Distress: Protocol for a Delphi Study

Sally Pezaro, DipMid, BA (Hons), RM, MSc; Wendy Clyne, BA (Hons), PhD

Centre for Technology Enabled Health Research, Faculty of Health and Life Sciences, Coventry University, Coventry, United Kingdom

Corresponding Author:

Sally Pezaro, DipMid, BA (Hons), RM, MSc
Centre for Technology Enabled Health Research
Faculty of Health and Life Sciences
Coventry University
Coventry, CV1 5FB
United Kingdom
Phone: 44 7950035977
Email: pezaros@uni.coventry.ac.uk

Abstract

Background: The development of an online intervention designed to effectively support midwives in work-related psychological distress will be challenging due to the ethical, practical, and therapeutic issues surrounding its design. Related literature suggests that midwives may require an anonymous, confidential, and therapeutic platform that facilitates amnesty and nonpunitive approaches to remedy ill health. However, it is unclear which requirements may be most salient to midwifery populations.

Objective: The objective of this paper is to describe the design of a Delphi study, intended to achieve expert consensus on the needs of midwives in work-related psychological distress who may be supported via an online intervention. This protocol may also serve as a research framework for similar studies to be modeled upon.

Methods: A heterogeneous sample of at least thirty experts on psychological well-being and distress associated with midwifery work will be recruited. Their opinions regarding the development of an online intervention designed to support midwives in work-related psychological distress will be collected through 2 rounds of questioning, via the Delphi Technique. When 60% (≥ 18 , assuming the minimum is 30) of panelists score within 2 adjacent points on a 7-point scale, consensus will be acknowledged. This Delphi study protocol will invite both qualitative and quantitative outcomes.

Results: This study is currently in development. It is financially supported by a full-time scholarship at the Centre for Technology Enabled Health Research at Coventry University (Coventry, UK). The implementation of this Delphi study is anticipated to occur during the autumn of 2015.

Conclusions: The results of this study will direct the development of an online intervention designed to support midwives in work-related psychological distress, summarize expert driven consensus, and direct future research.

(*JMIR Res Protoc* 2015;4(3):e107) doi: [10.2196/resprot.4766](https://doi.org/10.2196/resprot.4766)

KEYWORDS

Delphi technique; Internet; intervention studies; midwifery; psychological; research protocols; self-help groups; stress

Introduction

Background

The mental health and well-being of health care professionals has gathered significant attention due to its direct correlation with quality patient care [1]. Midwives may be at an increased risk of developing psychological distress due to the traumatic work environments they endure [2]. These environments report

incidents of workplace bullying, emotionally demanding clinical case loads, and a pressure to work despite feeling unwell enough to do so [1,2]. Interventions designed to support midwives in work-related psychological distress are required if the global shortage of midwives and the poor effects that midwives' psychological distress has on patient care are to be remedied. It is unclear who may be responsible for the well-being of health care staff in the United Kingdom, yet it is clear that there is a paucity of support for midwives in distress [3].

Midwives generally find it challenging to disclose personal experiences of psychological distress [4,5]. In addition, health care professionals who experience the distressing effects of functioning within traumatic work environments may not recognize mental ill health in themselves [6,7].

To enable midwives to seek help with the consequences of work-related psychological distress, a platform of amnesty, confidentiality, and anonymity may be required before any benefits may accrue [8,9].

Can Online Interventions Be the Answer?

An online intervention may be one solution that midwives may turn to in work-related psychological distress, as a preferred option of support [10]. To develop an online intervention that fits the needs of midwives, their employers, and professional bodies, it will be important to first define what characteristics an online intervention should have.

This paper outlines a protocol for a Delphi study designed to achieve expert consensus about what midwives in work-related psychological distress may need to be supported via an online intervention and peer support platform. The expert consensus will be used to inform the development and content of an online intervention for midwives in work-related psychological distress.

Methods

The Delphi Study Design

The Delphi Technique has been used extensively within health, social science, and intervention research [11-13]. It involves rounds of discussion whereby experts are invited to disclose their opinions on particular topics for which there is a paucity of knowledge. It is assumed that the opinions of many outweigh those of the individual, and thus, any consensus generated may be considered to be a valid expert opinion [14,15]. Because there is an incomplete state of knowledge about what midwives in work-related psychological distress may require when accessing an online intervention designed to effectively support them, a Delphi study was considered to be a suitable research tool to augment unanimity in opinion [16]. The distinct characteristics of the Delphi technique are (1) anonymity, (2) iteration, (3) controlled feedback, and (4) statistical “group response” [17].

Achieving consensus is the primary aim of the Delphi study, yet the measurement of consensus varies greatly [18]. There is no firm consensus as to what may be considered a consensus within a Delphi study. Within this Delphi study, a primary

criterion is that at least 60% (≥ 18 , assuming the minimum is 30) of Delphi panel members must indicate a preference within 2 adjacent response points on a 7-point Likert scale for consensus to be reached.

Rigid Delphi study designs have been criticized for their inability to allow their experts to elaborate on their opinions [12]. Therefore, this Delphi design will be a modified one [15,19]. Free text response options will accompany each statement put to panel members [20] to provide experts with the opportunity to elaborate on their opinions.

The research team who will conduct this Delphi study includes 6 academics with professional backgrounds in midwifery, general medicine, psychology, and academic research.

Participants

There are no clear guidelines in relation to what panel size is most appropriate for a Delphi study design [20]. A minimum of 30 experts will be recruited to this Delphi panel. Heterogeneity within the expert panel will play an essential part in ensuring study quality [16]. Therefore, panel members will be selected from different fields relating to midwifery care, health care, psychological distress, professional practice, and academia. They will be identified through a stakeholder analysis (see [Multimedia Appendix 1](#)). These experts will be midwives, researchers, lecturers, health care professionals, students, patient groups, and maternity-based organizations. Inclusion criteria are shown in [Figure 1](#).

Once experts have been identified, they will be directed toward information about the aim and content of the Delphi study. A formal invitation will also be given (see [Multimedia Appendix 2](#) and [21-31]). Potential participants will be invited to consent to participate as the online Delphi study begins (see [Multimedia Appendix 3](#)). Potential and recruited panel members will also be asked to refer other suitable individuals. This layer of recruitment aims to eliminate any bias from the research teams' recruitment selection. Solicitation of nominations of appropriate field experts is typically recommended as best practice in the Delphi study design [32].

Informed consent will be obtained from all participants as the first round of questioning begins online, and will include the consensual agreement to publish anonymized data and nonidentifiable data results (see [Multimedia Appendix 3](#)). Participants will be directed to appropriate support services both online and offline due to the sensitive nature of the subject matter. Participants will also receive copies of any publications that may result from the study and a summary of outcomes.

Figure 1. Participant inclusion criteria.

Expert Panel Inclusion Criteria
Either: A listed author in at least one publication relevant to
-Midwifery
-Psychology
-Psychological trauma
-Psychiatry
-Health care services
And/or: Practical knowledge in
-Midwifery
-Midwifery education
-Research
-Therapies
-Health services
-Patient experience
-Staff experience


Participant Recruitment

Overview

Experts will be invited to participate by the research team. They will be invited via email and social media contact with a formal invitation to become a part of the panel (see [Multimedia Appendix 2](#)). [Figure 2](#) shows the flowchart for participant recruitment.

It is anticipated that some experts may withdraw from the study during its course [33]. Therefore, social media will also be used to recruit participants to compensate for potential dropouts. A minimum of 30 panel members will be recruited to this study, although the team recognizes that there is no consensus regarding what the optimal number of participants for a Delphi study may be [14,34]. Should less than 50 experts be recruited before the Delphi study commences, an additional 50 people will be invited to participate to compensate for potential dropout rates and to avoid a failure to achieve adequate panel numbers.

Figure 2. Flowchart for participant recruitment.



Social Networking Recruitment

The research team will consult their social, academic, and occupational networks to identify potential experts who meet the inclusion criteria. Suitable candidates will receive an email inviting them to participate in the Delphi study.

Twitter will also be used for research recruitment due to its high-quality health care, research, and academic communities. Twitter is evidenced to be a highly effective tool for health care research recruitment [35]. Stakeholder groups identified in the

stakeholder analysis will then be asked to promote the study to their online followers. A link to a blog page with inclusion criteria, further information, support resources, and an online survey will be provided to facilitate online recruitment [36]. Willing and suitable participants can then express their interest in partaking in the study by contacting the research team directly.

Recruitment Through the Academic Literature

Experts within the field of midwifery, psychology, psychiatry, and health care will be identified through literature searching. The research team will identify key papers of relevance, the authors of which will then be invited to participate. They will be invited via email and social media contact with a formal invitation to become a part of the panel (see [Multimedia Appendix 2](#)).

Procedure

Overview

This Delphi study will employ the principles of anonymity, repetitions at each stage of questioning, and feedback between rounds of descriptive statistics regarding the group's response and summaries of free text responses about each item in the item panel [37]. The Delphi study technique was chosen as it prevents dominant individuals from controlling the process of group discussion [16]. This is particularly salient in hierarchal environments, such as the health care system, where many participants are anticipated to originate. The anonymity the Delphi study facilitates can also allow for unashamed freedom of speech, which in turn, leads to a more accurate opinion giving [38].

Experts will only be sent further correspondence should they indicate an initial interest to participate in the study. In the absence of any response to the initial invitation sent by the research team, it will be assumed that the recipient has no interest in participating in the study, and will therefore receive no further correspondence.

Experts who continue to participate within the study but do not respond to the first Delphi round will be sent 2 reminders via email or social media contact. To withdraw from the study, experts must directly contact the research team and explicitly state their withdrawal. Unless this action is confirmed, all experts will receive reminders and survey links for each round. Two weeks will be allocated for Delphi experts to respond to each round of questioning [39]. In total, there will be a 5-week interval between the initiation of the first round and the start of the second round of questioning.

Reminders will be sent to participants 1 week before each round begins in order to maximize their participation. A link to the survey will then be given to all participants.

Questions

Questions have been designed to explore consensus about the design, construction, purpose, and content of an online intervention to support midwives in work-related psychological distress (see [Multimedia Appendix 4](#)). These questions were developed in response to a review of the literature. This is an acceptable and a common modification of the Delphi process [39]. Literature reviewing remained broad in scope and included a combination of the search terms "burnout," "psychological distress," "midwives," "midwifery," "midwife," "online intervention," "self-help groups," "CBT," "mindfulness," "stress," "depression," "anxiety," "peer support," "mental health literacy," "second victim," "PTSD," "post-traumatic stress," "workplace bullying," and "NHS." In reading and re-reading

the retrieved literature, a theoretical basis was developed for what may or may not be useful in the development of an online intervention designed to support midwives in work-related psychological distress. These theories are put forward for testing before the expert panel.

There will be 3 themes of questioning and 2 response options available. The 3 themes will be intervention design and practical inclusions, inclusions of therapeutic support, and ethical inclusions. The 2 response options available will be a 7-Point Likert scale and open text responses.

Delphi Survey Design

Bristol Online Survey [40] will be used to administer the Delphi study. Round 1 will consist of a structured questionnaire. Respondents will be asked to indicate their priority rating for a series of items via Likert scale responses. They will also have the option to disclose why they chose to mark each item with lower or higher priority within an open text field. Respondents will also be invited to provide additional comments through the provision of a free text response. Finally, panelists will have the opportunity to suggest new questions to be put forward during the second round of questioning.

Round 2 will consist of a second questionnaire that is based on the information provided in the first round. The primary aim of this round will be to offer the panel the opportunity to reconsider their responses from Round 1 for those items for which consensus was not achieved in Round 1. This opportunity will be offered in light of feedback about the groups' responses in Round 1. New questions may also be added to this second round in response to suggestions put forward by the panel during the first round. Respondents will be asked to review these new questions and indicate their priority rating. Respondents will be invited to provide comments through the provision of a free text response option for each item in the second questionnaire. They will also again be given the opportunity to disclose why they have chosen to mark each item with lower or higher priority within an open text field.

Analysis

Because there are no conclusive guidelines for establishing consensus in Delphi literature [41], taking account of the average accord and the 7-point scale, consensus will be reached if 60% (≥ 18 , assuming the minimum is 30) of respondents are within 2 adjacent response points on the 7-point scale (eg, if 60%, ≥ 18 assuming the minimum is 30, of participants select 2 and 3 in response to a specific item). Items which do not achieve consensus in Round 1 will be re-presented in Round 2.

The mean, minimum, and maximum scores for each item will also be calculated and reported to panel members as feedback after each round.

Any free text responses provided by participants to specific items will be analyzed with thematic analysis [42]. Themes may be reframed, reviewed, and revised throughout this thematic analysis, as coherent patterns are formed. This thematic analysis of qualitative open responses will be presented in a table format and feedback will be provided to panel members after each round.

Results

This study is currently in development. It is financially supported by a full-time scholarship at the Centre for Technology Enabled Health Research at Coventry University (Coventry, UK). Ethical approval for this study has been granted by Coventry University Ethics Department. The implementation of this Delphi study is anticipated to occur during the autumn of 2015. Project reference id P35069.

Discussion

Preliminary Agenda

The aim of this Delphi study is to reach consensus on the salient themes and elements to be included within an online intervention to support midwives in work-related psychological distress. The results of this research will be used to inform the development of an online intervention designed to support midwives in psychological distress.

A key weakness of the Delphi technique is that it lacks a theoretical framework [14]. The advantage of using a Delphi

study technique within this research will be that ideas, definitions, and experiences of a variety of experts can be synthesized to inform development of the intervention. Panel members will be drawn from a variety of backgrounds, and as such will be able to contribute a variety of evidence and multidisciplinary perspectives.

Biases may occur in Delphi studies that could also distort the consensus. Desirability bias from both the experts and the research team could impede the achievement of a “true” consensus [43]. There is also the risk of ambiguity and conditional statements given within the questionnaire [44]. In this case, panel members may be interpreting the questions and statements differently. This may also lead to a polarization in results. A 7-point judgment scale is used to avoid elements of ambiguity; however, this may not protect results against some polarity [45]. To mitigate these risks, the questionnaire has been reviewed and piloted among peers.

Conclusions

This paper describes the design of a Delphi study. This will be the first Delphi study to explore the online support needs of midwives in work-related psychological distress.

Acknowledgments

This study is financially supported by a full-time scholarship, funded by the Centre for Technology Enabled Health Research at Coventry University.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Stakeholder analysis.

[\[PDF File \(Adobe PDF File\), 97KB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

Formal invitation presented to potential participants.

[\[PDF File \(Adobe PDF File\), 102KB-Multimedia Appendix 2\]](#)

Multimedia Appendix 3

Informed consent form.

[\[PDF File \(Adobe PDF File\), 92KB-Multimedia Appendix 3\]](#)

Multimedia Appendix 4

Delphi study questionnaire.

[\[PDF File \(Adobe PDF File\), 200KB-Multimedia Appendix 4\]](#)

References

1. Royal College of Physicians. Work and Well-Being in the NHS: Why Staff Health Matters to Patient Care. London, UK: Royal College of Physicians; 2015 Mar 13. URL: <https://www.rcplondon.ac.uk/sites/default/files/work-and-wellbeing-in-the-nhs.pdf> [accessed 2015-08-31] [WebCite Cache ID 6bBzaHVvo]
2. Hunter B, Warren L. Midwives experiences of workplace resilience. *Midwifery* 2014 Aug;30(8):926-934. [doi: [10.1016/j.midw.2014.03.010](https://doi.org/10.1016/j.midw.2014.03.010)] [Medline: [24742637](https://pubmed.ncbi.nlm.nih.gov/24742637/)]

3. Strobl J, Panesar S, Carson-Stevens A, McIldowie B, Ward H, Cross H, et al. Suicide by Clinicians Involved in Serious Incidents in the NHS: A Situational Analysis. Salford, UK: Clinical Leaders Network; 2014 Jun. URL: <http://www.cln.nhs.uk/userfiles/file/FINAL%20REPORT%20Suicides%20by%20clinicians%20involved%20in%20SIs%20in%20the%20NHS.pdf> [accessed 2015-08-31] [WebCite Cache ID 6bBzoN44U]
4. Ullström S, Sachs A, Hansson J, Øvretveit S, Brommels M. Suffering in silence: A qualitative study of second victims of adverse events. *BMJ Qual Saf* 2014 Apr;23(4):325-331. [doi: [10.1136/bmjqs-2013-002035](https://doi.org/10.1136/bmjqs-2013-002035)]
5. Sheen K, Spiby H, Slade P. Exposure to traumatic perinatal experiences and posttraumatic stress symptoms in midwives: Prevalence and association with burnout. *Int J Nurs Stud* 2015 Feb;52(2):578-587. [doi: [10.1016/j.ijnurstu.2014.11.006](https://doi.org/10.1016/j.ijnurstu.2014.11.006)] [Medline: [25561076](https://pubmed.ncbi.nlm.nih.gov/25561076/)]
6. Trippany R, Kress V, Wilcoxon S. Preventing vicarious trauma: What counsellors should know when working with trauma survivors. *J Couns Dev* 2004;82(1):31-37. [doi: [10.1002/j.1556-6678.2004.tb00283.x](https://doi.org/10.1002/j.1556-6678.2004.tb00283.x)]
7. Perry L, Lamont S, Brunero S, Gallagher R, Duffield C. The mental health of nurses in acute teaching hospital settings: A cross-sectional survey. *BMC Nurs* 2015;14:15 [FREE Full text] [doi: [10.1186/s12912-015-0068-8](https://doi.org/10.1186/s12912-015-0068-8)] [Medline: [25904820](https://pubmed.ncbi.nlm.nih.gov/25904820/)]
8. Munro R. Sick day scrutiny. *Nurs Stand* 2011;25(18):24-25. [Medline: [21309319](https://pubmed.ncbi.nlm.nih.gov/21309319/)]
9. Swami V. Mental health literacy of depression: Gender differences and attitudinal antecedents in a representative British sample. *PLoS One* 2012;7(11):e49779 [FREE Full text] [doi: [10.1371/journal.pone.0049779](https://doi.org/10.1371/journal.pone.0049779)] [Medline: [23166769](https://pubmed.ncbi.nlm.nih.gov/23166769/)]
10. Health Service Journal. Challenge Top-Down Change. London: EMAP Publishing Ltd; 2015. URL: <http://www.hs.j.co.uk/leadership/change-challenge> [accessed 2015-08-31] [WebCite Cache ID 6bC0g2nJO]
11. Efstathiou N, Ameen J, Coll A. A Delphi study to identify healthcare users' priorities for cancer care in Greece. *Eur J Oncol Nurs* 2008 Sep;12(4):362-371. [doi: [10.1016/j.ejon.2008.04.010](https://doi.org/10.1016/j.ejon.2008.04.010)] [Medline: [18547866](https://pubmed.ncbi.nlm.nih.gov/18547866/)]
12. Walker A, Selfe J. The Delphi method: A useful tool for the allied health researcher. *Br J Ther Rehab* 1996 Dec;3(12):677-681. [doi: [10.12968/bjtr.1996.3.12.14731](https://doi.org/10.12968/bjtr.1996.3.12.14731)]
13. Webber M, Reidy H, Ansari D, Stevens M, Morris D. Developing and modeling complex social interventions: Introducing the connecting people intervention. *Res Soc Work Pract* 2015 Mar 30 (forthcoming). [doi: [10.1177/1049731515578687](https://doi.org/10.1177/1049731515578687)]
14. Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs* 2000 Oct;32(4):1008-1015. [Medline: [11095242](https://pubmed.ncbi.nlm.nih.gov/11095242/)]
15. Habibi A, Sarafrazi A, Izadyar S. Delphi technique theoretical framework in qualitative research. *Int J Eng Sci* 2014;3(4):8-13.
16. Powell C. The Delphi technique: Myths and realities. *J Adv Nurs* 2003 Feb;41(4):376-382. [Medline: [12581103](https://pubmed.ncbi.nlm.nih.gov/12581103/)]
17. Rowe G, Wright G. Expert opinions in forecasting: The role of the Delphi technique. In: Armstrong JS, editor. *Principles of Forecasting: A Handbook for Researchers and Practitioners*. Boston, MA: Kluwer Academic Publishers; 2001:125-144.
18. von der Gracht HA. Consensus measurement in Delphi studies: Review and implications for future quality assurance. *Technol Forecast Soc Change* 2012 Oct;79(8):1525-1536.
19. Beretta R. A critical review of the Delphi technique. *Nur Res* 1996;3:79-89.
20. Keeney S, Hasson F, McKenna HP. A critical review of the Delphi technique as a research methodology for nursing. *Int J Nurs Stud* 2001 Apr;38(2):195-200. [Medline: [11223060](https://pubmed.ncbi.nlm.nih.gov/11223060/)]
21. Health and Safety Executive. *Stress-Related and Psychological Disorders in Great Britain 2014*. London, UK: Health and Safety Executive; 2014. URL: <http://www.hse.gov.uk/statistics/causdis/stress/index.htm> [accessed 2015-08-31] [WebCite Cache ID 6bC3Q9YKL]
22. Romani M, Ashkar K. Burnout among physicians. *Libyan J Med* 2014 Feb 17;9:23556. [doi: [10.3402/ljm.v9.23556](https://doi.org/10.3402/ljm.v9.23556)]
23. Rice H, Warland J. Bearing witness: Midwives experiences of witnessing traumatic birth. *Midwifery* 2013 Sep;29(9):1056-1063. [doi: [10.1016/j.midw.2012.12.003](https://doi.org/10.1016/j.midw.2012.12.003)] [Medline: [23415352](https://pubmed.ncbi.nlm.nih.gov/23415352/)]
24. McHugh A, Dawson L, Moncrieff G, McDonald B, Parton S, Gillies H. Are in You in Despair for Your Future in General Practice. 2014 Aug 15. URL: <https://pracmanhealth.files.wordpress.com/2014/08/are-you-in-despair-for-your-future-in-general-practice-final-report1.pdf> [accessed 2015-08-31] [WebCite Cache ID 6bC3oJK8o]
25. Royal College of Midwives. *High Quality Midwifery Care*. London, UK: Royal College of Midwives; 2014 Oct. URL: <https://www.rcm.org.uk/sites/default/files/High%20Quality%20Midwifery%20Care%20Final.pdf> [accessed 2015-08-31] [WebCite Cache ID 6bC3yw3D0]
26. NHS Litigation Authority. *Learning from Maternity Claims*. London, UK: NHS Litigation Authority; 2014. URL: <http://www.nhs.uk/CurrentActivity/Documents/Learning%20from%20Maternity%20Claims.pdf> [accessed 2015-08-31] [WebCite Cache ID 6bC48tFFW]
27. Boorman S. *NHS Health and Well-Being Review Interim Report*. London, UK: The Stationery Office; 2009. URL: http://webarchive.nationalarchives.gov.uk/20130107105354/http://www.dh.gov.uk/prod_consum_dh/groups/dh_digitalassets/documents/digitalasset/dh_108910.pdf [accessed 2015-09-02] [WebCite Cache ID 6bFLCEGq]
28. Black C. *Working for a Healthier Tomorrow*. London, UK: The Stationery Office; 2008.
29. Black C. Why healthcare organisations must look after their staff. *Nurs Manag (Harrow)* 2012 Oct;19(6):27-30. [doi: [10.7748/nm2012.10.19.6.27.c9319](https://doi.org/10.7748/nm2012.10.19.6.27.c9319)] [Medline: [23252089](https://pubmed.ncbi.nlm.nih.gov/23252089/)]
30. Department of Health. *Healthcare Sector Staff Wellbeing, Service Delivery, and Health Outcomes. A Compendium of Factsheets: Well-Being Across the Life Course*. London, UK: Department of Health; 2014 Jan. URL: <https://www.gov.uk/>

- [government/uploads/system/uploads/attachment_data/file/277591/Staff_wellbeing_service_delivery_and_health_outcomes.pdf](#) [accessed 2015-08-31] [WebCite Cache ID 6bC4VVEQS]
31. Ziebland S, Wyke S. Health and illness in a connected world: How might sharing experiences on the internet affect people's health? *Milbank Q* 2012 Jun;90(2):219-249 [FREE Full text] [doi: [10.1111/j.1468-0009.2012.00662.x](https://doi.org/10.1111/j.1468-0009.2012.00662.x)] [Medline: [22709387](https://pubmed.ncbi.nlm.nih.gov/22709387/)]
 32. Ludwig BG. Internationalizing Extension: An Exploration of the Characteristics Evident in a State University Extension System That Achieves Internationalization [dissertation]. Columbus, OH: The Ohio State University; 1994. URL: https://etd.ohiolink.edu/!etd.send_file?accession=osu1146146542&disposition=inline [accessed 2015-08-31]
 33. Evers S, Goossens M, de Vet H, van Tulder TM, Ament A. Criteria list for assessment of methodological quality of economic evaluations: Consensus on Health Economic Criteria. *Int J Technol Assess Health Care* 2005;21(2):240-245. [Medline: [15921065](https://pubmed.ncbi.nlm.nih.gov/15921065/)]
 34. Paré G, Cameron A, Poba-Nzaou P, Templier M. A systematic assessment of rigor in information systems ranking-type Delphi studies. *Inf Manage* 2013 Jul;50(5):207-217. [doi: [10.1016/j.im.2013.03.003](https://doi.org/10.1016/j.im.2013.03.003)]
 35. O'Connor A, Jackson L, Goldsmith L, Skirton H. Can I get a retweet please? Health research recruitment and the Twittersphere. *J Adv Nurs* 2014 Mar;70(3):599-609. [doi: [10.1111/jan.12222](https://doi.org/10.1111/jan.12222)] [Medline: [23909740](https://pubmed.ncbi.nlm.nih.gov/23909740/)]
 36. Pezaro S. Research Blog. 2015. URL: <https://healthystaff4healthypatients.wordpress.com/> [accessed 2015-08-31] [WebCite Cache ID 6bC249f9s]
 37. McKenna HP. The Delphi technique: A worthwhile research approach for nursing? *J Adv Nurs* 1994 Jun;19(6):1221-1225. [Medline: [7930104](https://pubmed.ncbi.nlm.nih.gov/7930104/)]
 38. Strauss H, Zeigler H. Delphi, political philosophy and the future. *Futures* 1975;7(3):184-196. [doi: [10.1016/0016-3287\(75\)90063-4](https://doi.org/10.1016/0016-3287(75)90063-4)]
 39. Hsu CC, Sandford BA. The Delphi technique: Making sense of consensus. *Pract Assess Res Eval* 2007;12(10):1-8.
 40. Bristol Online Survey. Bristol Online Survey 2015. 2015. URL: <http://www.survey.bris.ac.uk/> [accessed 2015-05-29] [WebCite Cache ID 6YsroN6XP]
 41. Clyne W, White S, McLachlan S. Developing consensus-based policy solutions for medicines adherence for Europe: A Delphi study. *BMC Health Serv Res* 2012 Dec 23;12:425 [FREE Full text] [doi: [10.1186/1472-6963-12-425](https://doi.org/10.1186/1472-6963-12-425)]
 42. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3(2):77-101.
 43. Ecken P, Gnatzy T, von der Gracht H. Desirability bias in foresight: Consequences for decision quality based on Delphi results. *Technol Forecast Soc Change* 2011 Nov;78(9):1654-1670. [doi: [10.1016/j.techfore.2011.05.006](https://doi.org/10.1016/j.techfore.2011.05.006)]
 44. Loveridge D. On Delphi Questions. Ideas in Progress. Manchester, UK: University of Manchester; 2002.
 45. Croasmun JT, Ostrom L. Using Likert-type scales in the social sciences. *J Adult Educ* 2011;40(1):19-22.

Edited by G Eysenbach; submitted 29.05.15; peer-reviewed by S McLachlan, D Menage; comments to author 15.07.15; revised version received 17.07.15; accepted 24.07.15; published 04.09.15

Please cite as:

Pezaro S, Clyne W

Achieving Consensus in the Development of an Online Intervention Designed to Effectively Support Midwives in Work-Related Psychological Distress: Protocol for a Delphi Study

JMIR Res Protoc 2015;4(3):e107

URL: <http://www.researchprotocols.org/2015/3/e107/>

doi: [10.2196/resprot.4766](https://doi.org/10.2196/resprot.4766)

PMID: [26341794](https://pubmed.ncbi.nlm.nih.gov/26341794/)

©Sally Pezaro, Wendy Clyne. Originally published in JMIR Research Protocols (<http://www.researchprotocols.org>), 04.09.2015. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Research Protocols, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.researchprotocols.org>, as well as this copyright and license information must be included.