

Protocol

Weight Loss After Stroke Through an Intensive Lifestyle Intervention (Group Lifestyle Balance-Cerebrovascular Accident): Protocol for a Randomized Controlled Trial

Simon Driver¹, PhD; Chad Swank¹, PT, PhD; Katherine Froehlich-Grobe¹, PhD; Evan McShan¹, MSc; Stephanie Calhoun¹, BSc; Monica Bennett², PhD

¹Baylor Scott and White Institute for Rehabilitation, Dallas, TX, United States

²Baylor Scott and White, Dallas, TX, United States

Corresponding Author:

Simon Driver, PhD

Baylor Scott and White Institute for Rehabilitation

909 N Washington Avenue

Dallas, TX, 75246

United States

Phone: 1 214 820 9014

Email: sjdriver@bswrehab.com

Abstract

Background: Weight gain can be a consequence of stroke, or cerebrovascular accident (CVA), because of impaired mobility, behavioral and emotional disorders, and sensory losses. Weight gain increases the patient's risk of recurrent stroke and chronic diseases, such as diabetes, metabolic syndrome, and pulmonary and heart disease. Approaches to weight loss in this population are lacking, although necessary because of the unique physiological and cognitive needs of persons after a stroke. Evidence shows that intensive behavioral therapy interventions that address both physical activity and diet offer the greatest potential for weight loss. The Group Lifestyle Balance (GLB) intervention is a 12-month, evidence-based weight loss program that has been used extensively with the general population; this program was modified to meet the needs of people who have had a stroke (GLB-CVA).

Objective: This randomized controlled trial (RCT) aims to examine the efficacy of the GLB-CVA on weight and secondary outcomes, compared with that of a waitlist control group.

Methods: This RCT will enroll and randomize 64 patients over an 18-month period.

Results: Currently, 51 people are waitlisted, with 23 out of 51 screened and 16 out of 23 eligible.

Conclusions: It is anticipated that the findings from this RCT will contribute to the evidence base regarding weight loss strategies for people living with stroke.

Clinical Trial: ClinicalTrials.gov NCT03873467; <https://clinicaltrials.gov/ct2/show/NCT03873467>

(*JMIR Res Protoc* 2019;8(10):e14338) doi: [10.2196/14338](https://doi.org/10.2196/14338)

KEYWORDS

cerebrovascular accident; physical activity; eating; weight loss; rehabilitation

Introduction

Background

Stroke, or cerebrovascular accident (CVA), is a serious public health issue because of its high incidence, high cost of care, and increased risk of morbidity and mortality [1]. Projections show that by 2030, an additional 3.4 million people aged >18 years will have had a stroke, which is a 20.5% increase in prevalence from 2012 [2]. Estimates indicate that about one-third of stroke

survivors have excessive weight, causing them to be overweight or obese (body mass index or BMI ≥ 25 kg/m²) [3]. Weight gain significantly restricts functioning and independence and greatly increases the risk of chronic diseases, such as diabetes, metabolic syndrome, pulmonary disease, heart disease, and recurrent stroke [4,5].

There is a high prevalence of weight gain and obesity after stroke, as people's physical activity and healthy eating behaviors are negatively impacted [3]. This may be because of a

combination of personal (eg, low motivation and lack of social support) and environmental factors (eg, lack of transportation, high cost of specialized programming, and lack of education from health care providers) [6]. Yet, increased physical activity and healthy dietary habits are recognized approaches to reduce the risk of comorbidities, such as obesity, hypertension, diabetes, recurrent stroke, and heart disease [7]. However, there is a lack of evidence-based approaches to weight loss for people after stroke [8]. Most currently available evidence-based weight loss programs were developed and tested with samples from the general population, with *disability* often used as an exclusion criterion [9,10]. Thus, the appropriateness of these programs to meet the unique needs of people who have had a stroke is unknown.

The Group Lifestyle Balance (GLB) program is a 12-month self-management intervention that has been shown to result in weight loss and reduce the risk for type 2 diabetes through improved physical activity and healthy eating behaviors [11-15]. Although the GLB has been used extensively with the general population [16,17], no study has been conducted specifically for people who have had a stroke. This protocol describes a randomized controlled trial (RCT) that will assess the efficacy of the GLB program that has been modified with input of stroke survivors (GLB-CVA). The GLB-CVA program will be compared with a waitlist control group. This protocol followed the standard protocol items: recommendations for interventional trials checklist to report relevant clinical trial details, as recommended by the Enhancing the Quality and Transparency of Health Research Network.

Objectives

Aim 1

Aim 1 of this study is to examine adherence to GLB-CVA intervention participation.

- *Hypothesis:* Intervention participants will attend at least 70% (15/22) of the weekly and monthly group-based sessions [18].

Aim 2

Aim 2 of this study is to conduct an RCT to examine the efficacy of the GLB-CVA on primary and secondary outcomes in the intervention group, compared with the waitlist control group at 3 and 6 months from baseline.


- *Hypothesis 2.1:* The intervention group will demonstrate statistically significant improvements in our primary (weight) and secondary outcomes (activity assessed with accelerometers, waist and arm circumference, blood pressure, hemoglobin A_{1c}, fasting glucose and lipid panel, 8-year diabetes risk, functional measures, self-report measures of physical activity and healthy eating, executive function, perceived social support, self-rated abilities for health behavior practices, pain interference, sleep disturbance, habits, and quality of life) when compared with the waitlist control group at 3 and 6 months.
- *Hypothesis 2.2:* Combined intervention data from both groups will demonstrate significant improvements in primary and secondary outcomes after 3, 6, and 12 months.

Methods

Study Design

The design is a single-phase, assessor-blinded, and waitlist-controlled RCT. Human subjects approval has been received by the Baylor Scott and White Research Institute Institutional Review Board (018-714); the study has been registered on ClinicalTrials.gov (NCT03873467). The intervention design and timeline are outlined in [Figure 1](#).

Figure 1. Enrollment and assessment schedule. GLB: Group Lifestyle Balance.



Study Setting

Study procedures will be completed at Baylor Scott and White Institute for Rehabilitation, a large rehabilitation hospital in an urban setting in the southwestern United States.

Participants and Recruitment

The primary sources of recruitment will be Baylor Scott and White Institute for Rehabilitation and community agencies in the Dallas–Fort Worth metroplex. Recruitment will be completed by trained interventionists who are members of the research team, and approaches include in-person visits to community organizations and stroke support groups, flyers, calls, and physician referrals, with a snowball technique for

further recruitment. Recruitment will occur over a 4- to 6-month period to ensure that the target sample size is reached.

Eligibility

Interested participants will contact or be contacted by the research team using information provided on approved study flyers. An eligibility screener will be completed telephonically with the following inclusion criteria: (1) BMI ≥ 25 kg/m², (2) aged 18 to 85 years, (3) all types of stroke, (4) at least 12 months after stroke, and (5) physician approval to be physically active and make dietary changes. Exclusion criteria are as follows: (1) low cognitive function (<10 on the Cognistat) [19], (2) residing in a hospital, acute rehab setting, or skilled nursing facility, (3) preexisting diagnosis of an eating disorder, (4) pregnancy, (5)

taking medication for diabetes, or (6) not fluent in English language. Before the completion of study procedures, informed consent will be obtained by trained research personnel in a private room at Baylor Scott and White Institute for Rehabilitation.

Intervention

The GLB intervention promotes self-management of a healthy lifestyle based on the tenets of Social Cognitive Theory [20] and the Health Belief Model [21]. The GLB program facilitates individual engagement in health behavior change and is completed in a group setting [11-15]. Data from previous GLB trials have shown 5% to 7% weight loss in a variety of settings, including community centers, churches, worksites, and health care systems [13,16,22-28]. The goal for participants is to reach 150 min of physical activity following the American Heart Association's (AHA) recommendations and follow dietary guidelines by the United States Department of Agriculture. The GLB program comprises 22 sessions over 12 months, including 12 weekly core program sessions, 4 transition phase sessions, and 6 support sessions (see [Multimedia Appendix 1](#)). GLB program materials are made available to the public under the Creative Commons licensing agreement through the Diabetes and Prevention Support Center at the University of Pittsburgh that developed the GLB program.

The GLB curriculum was adapted based on the recommendations from an advisory board of 29 key stakeholders, including stroke survivors, caregivers, clinicians (eg, physicians, therapists, registered dietitians, and neuropsychologists), community partners (eg, AHA and support groups), and researchers, with specific consideration for individuals with stroke (GLB-CVA). The advisory board met to review and discuss relevant changes to the existing curriculum during a 1-day meeting and were compensated US \$100 for their time. The recommended modifications were then made by the study team and approved by the Diabetes and Prevention Support Center at the University of Pittsburgh. Modifications included (1) reorganizing and refocusing the content to reflect the importance of healthy behaviors on heart health and prevent recurrent stroke, (2) reducing the volume of content to focus on 2 to 3 main points at each session, (3) involving care partner in the sessions to provide physical and emotional support, (4) developing stroke-specific handouts on weight loss barriers and healthy lifestyle importance, (5) creating and locating handouts and weblinks to modify physical activity and adaptive cooking, and (6) including guest lectures by experts in the field (eg, physical therapist and dietitian). The GLB-CVA intervention will be delivered by trained GLB lifestyle coaches at Baylor Scott and White Institute for Rehabilitation. Training was completed at either the University of Pittsburgh's Lifestyle Coach Training Workshop or Master Training Center at Baylor Scott and White Health and Wellness Institute. Both sites have recognition by the Centers for Disease Control and Prevention (CDC) as official GLB training centers.

Outcome Measures

Participants will complete assessments at 4 time points over the course of the yearlong intervention, including baseline, 3, 6, and 12 months. The waitlist control group will complete 2

additional assessment periods that serve as data for comparison in the RCT before they undergo the 12-month intervention. Demographic data collected will include the following: type of stroke, severity of disability (Modified Rankin Scale [29]), current age, date of stroke, sex, parental history of diabetes, race and ethnicity, education level, premorbid history of mental illness, marital and relationship status, diagnosed medical conditions, previous and present smoking and cigarettes per day, alcohol consumption and drinks per week, residence status and zip code, annual household income category, pre- and current stroke insurance type, pre- and current employment status, prestroke and current weight, height, and resting metabolic rate with MedGem. The primary and secondary outcomes are outlined in [Multimedia Appendix 2](#) [30-49]. Participants will be compensated US \$25 for their participation at each study time point. Assessments will be completed by a data collector who is trained as a medical assistant and phlebotomist and has completed Modified Rankin Scale training for previous stroke studies.

Sample Size

For the proposed study, 64 participants (32 per group) will be enrolled. The sample size was calculated to ensure that we have sufficient power for our analyses on the primary outcome (weight). Previous GLB studies with the general population have demonstrated weight loss over time (median $d=0.99$ at 3 months, 0.91 at 6 months, and 0.74 at 12 months) [15-18]. Results also indicated a maximum attrition rate of 16%, 23%, and 33% at 3, 6, and 12 months, respectively. Thus, a sample size of 64 will allow a power of greater than 90% to detect a conservatively assumed group difference of 4% in weight change (95% power). In the event of 33% attrition, we will achieve 83% to 87% power to detect these differences.

Allocation

Participants will be assigned to the experimental GLB-CVA or waitlist control groups using computer-generated random numbers in Microsoft Excel. Individuals in the waitlist control will be placed on a waiting list and receive the intervention after a 6-month period. To ensure an equal distribution between groups, block randomization will be used with blocks of sizes 4 and 6. Randomly mixing block sizes will reduce the study coordinator's ability to predict the last assignment of each block. The randomization list will be generated by the biostatistician, with results contained in sealed envelopes labeled with study identification numbers. After a participant is enrolled in the study, a study coordinator will select the assigned envelope to reveal the participant's group. Owing to the type of intervention, it is not practical to blind study participants to group assignment. However, to minimize assessor bias, outcome assessments will be conducted by a coordinator who (1) is blinded to group assignment, (2) is not included in study team meetings, (3) has a script to remind participants at the beginning of each assessment to maintain blinding, and (4) has a process for recording unblinding.

Data Management, Quality Assurance, and Exclusion of Bias

All nonelectronic source documents will be kept in a locked storage cabinet in the Baylor Scott and White Institute for Rehabilitation research office. Case report forms and all outcomes data will be inputted into REDCap (Vanderbilt, TN), a Health Insurance Portability and Accountability Act–compliant (21 Code of Federal Regulation Part 11) secure Web-based program, by trained study staff. All electronic data will be stored and maintained on a secure server and disposed in accordance with current federal guidelines.

Data management activities will occur quarterly and include data quality checks and verification, as well as internal logic checks (eg, outlier values and internal inconsistencies). Of the participant files, 10% will be audited for source document and data entry review. Cross-tabulation checks using SAS will also be applied to the data. Data will be stored and backed up periodically by the biostatistician on the secure server. Descriptive statistics will be prepared and included into quarterly reports to ensure the quality of data and study progress. The principal investigator will provide oversight on all data entry and proper data monitoring and audit procedures.

Statistical Methods

To examine adherence to the GLB-CVA intervention (Aim 1), univariate analysis will be used to summarize session attendance to determine if it reaches the hypothesized rate of at least 70% (15/22). An RCT will be completed to examine the efficacy of the GLB-CVA on primary and secondary outcomes in the intervention group compared with the waitlist control group at 3, 6, and 12 months from baseline. General mixed modeling analysis will be conducted for the primary and secondary outcomes to assess initial and sustained impacts of the adapted GLB-CVA intervention. More specifically, individual growth models will be evaluated for linear and nonlinear change from baseline to 3, 6, and 12 months (level 1, time effects), overall group difference (level 2, group effect), and group difference in change (cross-level, time-by-group interaction effect; Hypothesis 2.1). Models will be adjusted for demographic variables (eg, age, gender, ethnicity, and disability severity using the Modified Rankin Scale), particularly if they are imbalanced after randomization, thereby providing more accurate estimates of the intervention impacts.

Separate growth models for change from baseline to 12 months will be fitted within both groups combined (time effects only; Hypothesis 2.2). Data will include missing observations resulting from either attrition or nonresponse. Intent-to-treat analysis will use restricted maximum likelihood estimation, which can produce unbiased estimates with incomplete data. In addition, sensitivity analysis will utilize iterative Monte Carlo Markov Chain multiple imputation for missing data. All selected variables will be used in the imputation process, allowing for greater recovery of the missing data. All analyses will be performed using SAS 9.4.

Results

We currently have 51 people on the waitlist. Of these, 23 have been screened, and 16 of the 23 are eligible and scheduled for baseline assessments. The intervention is expected to start in the summer of 2019.

Discussion

The aim of our study is to examine the efficacy of the GLB-CVA on the weight as well as health and function of people who have had a stroke. As the fourth leading cause of disease burden globally, stroke and the resultant comorbidities (ie, hypertension, obesity, and diabetes) present a significant public health challenge, and interventions are therefore needed [50]. A combination of health behaviors (physical activity, nonsmoking, moderate alcohol intake, and adequate vitamin C intake) is known to reduce the incidence of stroke 2-fold [51]. Moreover, any lifestyle intervention attempting to improve health in people with stroke should include, as a primary focus, management of weight, blood pressure, cholesterol, and glucose [52]. Nonpharmacological interventions (ie, physical activity and healthy eating) are necessary, in addition to medication, to address these modifiable risk factors. However, a recent scoping review of the physical activity and stroke literature [53] found only 3 lifestyle intervention trials [54–56] that incorporated both physical activity and healthy eating strategies. Although the studies had short-term benefits (9–24 weeks) on mobility, dietary behaviors (salt intake), and metabolic factors (blood pressure and cholesterol), the longitudinal effects on weight, health, and function were not documented. As such, the GLB program was chosen because of data consistently demonstrating 5% to 7% weight loss, evidence of success in other disability populations [57] and programmatic flexibility to address the unique health and function needs of people post-CVA.

Although physical activity and healthy eating are foundational to evidence-based weight loss programs and recognized to reduce the risk of hypertension, obesity, diabetes, and heart disease [58–60], after stroke, people encounter significant challenges to maintaining a healthy lifestyle. Common barriers following stroke include personal (eg, low motivation and physical impairment) and environmental factors (eg, lack of accessible facility and no transport) [6,61–64]. These barriers, in addition to others (eg, lack of social support and limited financial resources), reduce the ability of people who have had a stroke to participate in community-based health and wellness programs, further increasing their risk of developing comorbidities (eg, obesity, diabetes, and heart disease). To adapt a healthy lifestyle intervention to meet the unique needs of people who have had a stroke, modifications must account for common barriers to healthy lifestyle behaviors and address frequent health risks after stroke.

The GLB-CVA program was modified from the original evidence-based curriculum, with inputs from 29 advisory board members, to meet the specific needs of people with stroke. This modified curriculum addresses the unique needs of people with stroke by incorporating the inclusion of care partners, stroke-specific handouts on weight loss barriers, reduced content

to 2 to 3 main points for each session, reorganization and refocus on the importance of healthy lifestyle on stroke prevention and heart health, weblinks, and guest lecturers for stroke-specific modifications to physical activity and adaptive cooking. Findings from this RCT are expected to establish a strong evidence-based approach to weight loss among people after stroke that is scalable into community settings throughout the nation.

Upon completion of this RCT, all GLB-CVA materials will be made accessible to the public at no charge through the Diabetes

Prevention and Support Center at the University of Pittsburgh. The GLB program was selected because of the fact that it is acknowledged by the CDC-National Diabetes Prevention Recognition Program (CDC-DPRP) as an evidence-based lifestyle program. In addition, Centers for Medicare and Medicaid Services recently included the GLB into the Medicare payment program, allowing individuals to be reimbursed for participation [65]. The primary goal of this RCT is to attain full recognition by the CDC-DPRP and achieve reimbursement from the Centers for Medicare and Medicaid Services for people with stroke living in the community.

Acknowledgments

The contents of this paper were developed under a grant from the National Institute on Disability, Independent Living, and Rehabilitation Research (NIDILRR grant number 90IFRE0021). NIDILRR is a center within the Administration for Community Living (ACL), Department of Health and Human Services (HHS). The contents of this paper do not necessarily represent the policy of NIDILRR, ACL, or HHS, and endorsement by the federal government should not be assumed.

Authors' Contributions

SD was responsible for study inception, design, oversight, and the first draft of the paper. CS and KFG contributed to the scientific merit and design of the study. MB provided sample size calculations and statistical support. EM and SC established study operations and recruitment. All authors have read and contributed to this paper.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Group Lifestyle Balance curriculum.

[\[PDF File \(Adobe PDF File\)137 KB-Multimedia Appendix 1\]](#)

Multimedia Appendix 2

Outcome measures for Group Lifestyle Balance-Cerebrovascular Accident project.

[\[PDF File \(Adobe PDF File\)108 KB-Multimedia Appendix 2\]](#)

References

1. O'Brien SR, Xue Y, Ingersoll G, Kelly A. Shorter length of stay is associated with worse functional outcomes for medicare beneficiaries with stroke. *Phys Ther* 2013 Dec;93(12):1592-1602. [doi: [10.2522/ptj.20120484](https://doi.org/10.2522/ptj.20120484)] [Medline: [23886846](https://pubmed.ncbi.nlm.nih.gov/23886846/)]
2. Ovbiagele B, Goldstein LB, Higashida RT, Howard VJ, Johnston SC, Khavjou OA, American Heart Association Advocacy Coordinating Committee and Stroke Council. Forecasting the future of stroke in the United States: a policy statement from the American Heart Association and American Stroke Association. *Stroke* 2013 Aug;44(8):2361-2375. [doi: [10.1161/STR.0b013e31829734f2](https://doi.org/10.1161/STR.0b013e31829734f2)] [Medline: [23697546](https://pubmed.ncbi.nlm.nih.gov/23697546/)]
3. Vahlberg B, Zetterberg L, Lindmark B, Hellström K, Cederholm T. Functional performance, nutritional status, and body composition in ambulant community-dwelling individuals 1-3 years after suffering from a cerebral infarction or intracerebral bleeding. *BMC Geriatr* 2016 Feb 19;16:48 [FREE Full text] [doi: [10.1186/s12877-016-0226-1](https://doi.org/10.1186/s12877-016-0226-1)] [Medline: [26895855](https://pubmed.ncbi.nlm.nih.gov/26895855/)]
4. Global Burden of Metabolic Risk Factors for Chronic Diseases Collaboration (BMI Mediated Effects), Lu Y, Hajifathalian K, Ezzati M, Woodward M, Rimm EB, et al. Metabolic mediators of the effects of body-mass index, overweight, and obesity on coronary heart disease and stroke: a pooled analysis of 97 prospective cohorts with 1.8 million participants. *Lancet* 2014 Mar 15;383(9921):970-983 [FREE Full text] [doi: [10.1016/S0140-6736\(13\)61836-X](https://doi.org/10.1016/S0140-6736(13)61836-X)] [Medline: [24269108](https://pubmed.ncbi.nlm.nih.gov/24269108/)]
5. Froehlich-Grobe K, Lollar D. Obesity and disability: time to act. *Am J Prev Med* 2011 Nov;41(5):541-545 [FREE Full text] [doi: [10.1016/j.amepre.2011.07.015](https://doi.org/10.1016/j.amepre.2011.07.015)] [Medline: [22011427](https://pubmed.ncbi.nlm.nih.gov/22011427/)]
6. Rimmer JH, Wang E, Smith D. Barriers associated with exercise and community access for individuals with stroke. *J Rehabil Res Dev* 2008;45(2):315-322 [FREE Full text] [doi: [10.1682/JRRD.2007.02.0042](https://doi.org/10.1682/JRRD.2007.02.0042)] [Medline: [18566948](https://pubmed.ncbi.nlm.nih.gov/18566948/)]
7. National Academies of Sciences, Engineering, and Medicine. 2012. Accelerating Progress in Obesity Prevention: Solving the Weight of the Nation URL: <http://www.nationalacademies.org/hmd/Reports/2012/Accelerating-Progress-in-Obesity-Prevention.aspx> [accessed 2017-05-03]

8. Jones TM, Dean CM, Hush JM, Dear BF, Titov N. A systematic review of the efficacy of self-management programs for increasing physical activity in community-dwelling adults with acquired brain injury (ABI). *Syst Rev* 2015 Apr 19;4:51 [FREE Full text] [doi: [10.1186/s13643-015-0039-x](https://doi.org/10.1186/s13643-015-0039-x)] [Medline: [25927591](https://pubmed.ncbi.nlm.nih.gov/25927591/)]
9. Tomlinson M, Swartz L, Officer A, Chan KY, Rudan I, Saxena S. Research priorities for health of people with disabilities: an expert opinion exercise. *Lancet* 2009 Nov 28;374(9704):1857-1862. [doi: [10.1016/S0140-6736\(09\)61910-3](https://doi.org/10.1016/S0140-6736(09)61910-3)] [Medline: [19944866](https://pubmed.ncbi.nlm.nih.gov/19944866/)]
10. Rimmer JH, Vanderbom KA, Bandini LG, Drum CE, Luken K, Suarez-Balcazar Y, et al. GRAIDs: a framework for closing the gap in the availability of health promotion programs and interventions for people with disabilities. *Implement Sci* 2014 Aug 14;9:100 [FREE Full text] [doi: [10.1186/s13012-014-0100-5](https://doi.org/10.1186/s13012-014-0100-5)] [Medline: [25123550](https://pubmed.ncbi.nlm.nih.gov/25123550/)]
11. Diabetes Prevention Program (DPP) Research Group. The Diabetes Prevention Program (DPP): description of lifestyle intervention. *Diabetes Care* 2002 Dec;25(12):2165-2171 [FREE Full text] [doi: [10.2337/diacare.25.12.2165](https://doi.org/10.2337/diacare.25.12.2165)] [Medline: [12453955](https://pubmed.ncbi.nlm.nih.gov/12453955/)]
12. Kramer MK, McWilliams JR, Chen H, Siminerio LM. A community-based diabetes prevention program: evaluation of the group lifestyle balance program delivered by diabetes educators. *Diabetes Educ* 2011;37(5):659-668. [doi: [10.1177/0145721711411930](https://doi.org/10.1177/0145721711411930)] [Medline: [21918204](https://pubmed.ncbi.nlm.nih.gov/21918204/)]
13. Kramer MK, Kriska AM, Venditti EM, Miller RG, Brooks MM, Burke LE, et al. Translating the diabetes prevention program: a comprehensive model for prevention training and program delivery. *Am J Prev Med* 2009 Dec;37(6):505-511. [doi: [10.1016/j.amepre.2009.07.020](https://doi.org/10.1016/j.amepre.2009.07.020)] [Medline: [19944916](https://pubmed.ncbi.nlm.nih.gov/19944916/)]
14. Kramer MK, Molenaar DM, Arena VC, Venditti EM, Meehan RJ, Miller RG, et al. Improving employee health: evaluation of a worksite lifestyle change program to decrease risk factors for diabetes and cardiovascular disease. *J Occup Environ Med* 2015 Mar;57(3):284-291 [FREE Full text] [doi: [10.1097/JOM.0000000000000350](https://doi.org/10.1097/JOM.0000000000000350)] [Medline: [25742535](https://pubmed.ncbi.nlm.nih.gov/25742535/)]
15. Ma J, Yank V, Xiao L, Lavori PW, Wilson SR, Rosas LG, et al. Translating the diabetes prevention program lifestyle intervention for weight loss into primary care: a randomized trial. *JAMA Intern Med* 2013 Jan 28;173(2):113-121 [FREE Full text] [doi: [10.1001/2013.jamainternmed.987](https://doi.org/10.1001/2013.jamainternmed.987)] [Medline: [23229846](https://pubmed.ncbi.nlm.nih.gov/23229846/)]
16. Kramer MK, Kriska AM, Venditti EM, Semler LN, Miller RG, McDonald T, et al. A novel approach to diabetes prevention: evaluation of the Group Lifestyle Balance program delivered via DVD. *Diabetes Res Clin Pract* 2010 Dec;90(3):e60-e63. [doi: [10.1016/j.diabres.2010.08.013](https://doi.org/10.1016/j.diabres.2010.08.013)] [Medline: [20863586](https://pubmed.ncbi.nlm.nih.gov/20863586/)]
17. Greenwood DA, Kramer MK, Hankins AI, Parise CA, Fox A, Buss KA. Adapting the Group Lifestyle Balance™ Program for weight management within a large health care system diabetes education program. *Diabetes Educ* 2014 May;40(3):299-307. [doi: [10.1177/0145721714524281](https://doi.org/10.1177/0145721714524281)] [Medline: [24562606](https://pubmed.ncbi.nlm.nih.gov/24562606/)]
18. Driver S, Reynolds M, Woolsey A, Callender L, Prajapati PK, Bennett M, et al. Impact of a community-based healthy lifestyle program on individuals with traumatic brain injury. *J Head Trauma Rehabil* 2018;33(6):E49-E58. [doi: [10.1097/HTR.0000000000000372](https://doi.org/10.1097/HTR.0000000000000372)] [Medline: [29385008](https://pubmed.ncbi.nlm.nih.gov/29385008/)]
19. Kiernan RJ, Mueller J, Langston W, van Dyke C. The Neurobehavioral Cognitive Status Examination: a brief but quantitative approach to cognitive assessment. *Ann Intern Med* 1987 Oct;107(4):481-485. [doi: [10.7326/0003-4819-107-4-481](https://doi.org/10.7326/0003-4819-107-4-481)] [Medline: [3631786](https://pubmed.ncbi.nlm.nih.gov/3631786/)]
20. Bandura A. Human agency in social cognitive theory. *Am Psychol* 1989 Sep;44(9):1175-1184. [doi: [10.1037/0003-066X.44.9.1175](https://doi.org/10.1037/0003-066X.44.9.1175)] [Medline: [2782727](https://pubmed.ncbi.nlm.nih.gov/2782727/)]
21. Hochbaum G, Rosenstock I, Kegels S. Academia.edu. 1952. Health Belief Model URL: http://www.academia.edu/download/37641514/Health_Belief_Model.doc [accessed 2019-08-08]
22. Kramer SF, Cumming T, Churilov L, Bernhardt J. Measuring activity levels at an acute stroke ward: comparing observations to a device. *Biomed Res Int* 2013;2013:460482 [FREE Full text] [doi: [10.1155/2013/460482](https://doi.org/10.1155/2013/460482)] [Medline: [24282815](https://pubmed.ncbi.nlm.nih.gov/24282815/)]
23. Davis-Smith YM, Davis-Smith M, Boltri JM, Seale JP, Shellenberger S, Blalock T, et al. Implementing a diabetes prevention program in a rural African-American church. *J Natl Med Assoc* 2007 Apr;99(4):440-446. [Medline: [17444435](https://pubmed.ncbi.nlm.nih.gov/17444435/)]
24. Amundson HA, Butcher M, Gohdes D, Hall T, Harwell TS, Helgerson SD, Montana Cardiovascular Disease and Diabetes Prevention Program Workgroup. Translating the diabetes prevention program into practice in the general community: findings from the Montana cardiovascular disease and diabetes prevention program. *Diabetes Educ* 2009;35(2):209-10,213. [doi: [10.1177/0145721709333269](https://doi.org/10.1177/0145721709333269)] [Medline: [19321807](https://pubmed.ncbi.nlm.nih.gov/19321807/)]
25. Ackermann RT, Finch EA, Brizendine E, Zhou H, Marrero DG. Translating the diabetes prevention program into the community. The DEPLOY pilot study. *Am J Prev Med* 2008 Oct;35(4):357-363 [FREE Full text] [doi: [10.1016/j.amepre.2008.06.035](https://doi.org/10.1016/j.amepre.2008.06.035)] [Medline: [18779029](https://pubmed.ncbi.nlm.nih.gov/18779029/)]
26. Vadheim LM, Brewer KA, Kassner DR, Vanderwood KK, Hall TO, Butcher MK, et al. Effectiveness of a lifestyle intervention program among persons at high risk for cardiovascular disease and diabetes in a rural community. *J Rural Health* 2010;26(3):266-272. [doi: [10.1111/j.1748-0361.2010.00288.x](https://doi.org/10.1111/j.1748-0361.2010.00288.x)] [Medline: [20633095](https://pubmed.ncbi.nlm.nih.gov/20633095/)]
27. Katula JA, Vitolins MZ, Morgan TM, Lawlor MS, Blackwell CS, Isom SP, et al. The healthy living partnerships to prevent diabetes study: 2-year outcomes of a randomized controlled trial. *Am J Prev Med* 2013 Apr;44(4 Suppl 4):S324-S332. [doi: [10.1016/j.amepre.2012.12.015](https://doi.org/10.1016/j.amepre.2012.12.015)] [Medline: [23498294](https://pubmed.ncbi.nlm.nih.gov/23498294/)]

28. Katula JA, Vitolins MZ, Rosenberger EL, Blackwell CS, Morgan TM, Lawlor MS, et al. One-year results of a community-based translation of the Diabetes Prevention Program: healthy-living partnerships to prevent diabetes (HELP PD) project. *Diabetes Care* 2011 Jul;34(7):1451-1457 [FREE Full text] [doi: [10.2337/dc10-2115](https://doi.org/10.2337/dc10-2115)] [Medline: [21593290](https://pubmed.ncbi.nlm.nih.gov/21593290/)]
29. Dromerick AW, Edwards DF, Diringner MN. Sensitivity to changes in disability after stroke: a comparison of four scales useful in clinical trials. *J Rehabil Res Dev* 2003;40(1):1-8 [FREE Full text] [doi: [10.1682/JRRD.2003.01.0001](https://doi.org/10.1682/JRRD.2003.01.0001)] [Medline: [15150715](https://pubmed.ncbi.nlm.nih.gov/15150715/)]
30. Compagnat M, Mandigout S, Chaparro DA, Daviet JC, Salle JY. Validity of the Actigraph GT3x and influence of the sensor positioning for the assessment of active energy expenditure during four activities of daily living in stroke subjects. *Clin Rehabil* 2018 Dec;32(12):1696-1704. [doi: [10.1177/0269215518788116](https://doi.org/10.1177/0269215518788116)] [Medline: [30012036](https://pubmed.ncbi.nlm.nih.gov/30012036/)]
31. Sirichana W, Dolezal BA, Neufeld EV, Wang X, Cooper CB. Wrist-worn triaxial accelerometry predicts the energy expenditure of non-vigorous daily physical activities. *J Sci Med Sport* 2017 Aug;20(8):761-765. [doi: [10.1016/j.jsams.2017.01.233](https://doi.org/10.1016/j.jsams.2017.01.233)] [Medline: [28159535](https://pubmed.ncbi.nlm.nih.gov/28159535/)]
32. Gersten JW, Orr W. External work of walking in hemiparetic patients. *Scand J Rehabil Med* 1971;3(1):85-88. [Medline: [5156160](https://pubmed.ncbi.nlm.nih.gov/5156160/)]
33. Serra MC, Balraj E, DiSanzo BL, Ivey FM, Hafer-Macko CE, Treuth MS, et al. Validating accelerometry as a measure of physical activity and energy expenditure in chronic stroke. *Top Stroke Rehabil* 2017 Jan;24(1):18-23 [FREE Full text] [doi: [10.1080/10749357.2016.1183866](https://doi.org/10.1080/10749357.2016.1183866)] [Medline: [27322733](https://pubmed.ncbi.nlm.nih.gov/27322733/)]
34. Palmer-McLean K, Harbst K. Stroke and brain injury. In: Moore G, Painter P, Roberts S, editors. *ACSM's Exercise Management for Persons with Chronic Diseases and Disabilities*. Champaign, IL: Human Kinetics; 2009:238-246.
35. Middleton A, Fritz SL, Lusardi M. Walking speed: the functional vital sign. *J Aging Phys Act* 2015 Apr;23(2):314-322 [FREE Full text] [doi: [10.1123/japa.2013-0236](https://doi.org/10.1123/japa.2013-0236)] [Medline: [24812254](https://pubmed.ncbi.nlm.nih.gov/24812254/)]
36. Wilson PW, Meigs JB, Sullivan L, Fox CS, Nathan DM, D'Agostino RB. Prediction of incident diabetes mellitus in middle-aged adults: the Framingham offspring study. *Arch Intern Med* 2007 May 28;167(10):1068-1074. [doi: [10.1001/archinte.167.10.1068](https://doi.org/10.1001/archinte.167.10.1068)] [Medline: [17533210](https://pubmed.ncbi.nlm.nih.gov/17533210/)]
37. Block G, Azar KM, Romanelli RJ, Block TJ, Hopkins D, Carpenter JA, et al. Diabetes prevention and weight loss with a fully automated behavioral intervention by email, web, and mobile phone: a randomized controlled trial among persons with prediabetes. *J Med Internet Res* 2015 Oct 23;17(10):e240 [FREE Full text] [doi: [10.2196/jmir.4897](https://doi.org/10.2196/jmir.4897)] [Medline: [26499966](https://pubmed.ncbi.nlm.nih.gov/26499966/)]
38. Delacre M, Lakens D, Leys C. Why psychologists should by default use Welch's t-test instead of student's t-test. *Int Rev Soc Psychol* 2017 Apr 5;30(1):92-101 [FREE Full text] [doi: [10.5334/irsp.82](https://doi.org/10.5334/irsp.82)]
39. Salter KL, Moses MB, Foley NC, Teasell R. Health-related quality of life after stroke: what are we measuring? *Int J Rehabil Res* 2008 Jun;31(2):111-117. [doi: [10.1097/MRR.0b013e3282fc0f33](https://doi.org/10.1097/MRR.0b013e3282fc0f33)] [Medline: [18467925](https://pubmed.ncbi.nlm.nih.gov/18467925/)]
40. Duncan PW, Bode RK, Lai SM, Perera S, Glycine Antagonist in Neuroprotection Americans Investigators. Rasch analysis of a new stroke-specific outcome scale: the stroke impact scale. *Arch Phys Med Rehabil* 2003 Jul;84(7):950-963. [doi: [10.1016/S0003-9993\(03\)00035-2](https://doi.org/10.1016/S0003-9993(03)00035-2)] [Medline: [12881816](https://pubmed.ncbi.nlm.nih.gov/12881816/)]
41. Brown SC, Pantin H, Lombard J, Toro M, Huang S, Plater-Zyberk E, et al. Walk Score®: associations with purposive walking in recent Cuban immigrants. *Am J Prev Med* 2013 Aug;45(2):202-206 [FREE Full text] [doi: [10.1016/j.amepre.2013.03.021](https://doi.org/10.1016/j.amepre.2013.03.021)] [Medline: [23867028](https://pubmed.ncbi.nlm.nih.gov/23867028/)]
42. Becker H, Stuifbergen A, Oh HS, Hall S. Self-rated abilities for health practices: a health self-efficacy measure. *Health Values* 1993;17(5):42-50 [FREE Full text]
43. Zimet GD, Dahlem NW, Zimet SG, Farley GK. The multidimensional scale of perceived social support. *J Pers Assess* 1988 Mar;52(1):30-41. [doi: [10.1207/s15327752jpa5201_2](https://doi.org/10.1207/s15327752jpa5201_2)]
44. Amtmann D, Cook KF, Jensen MP, Chen WH, Choi S, Revicki D, et al. Development of a PROMIS item bank to measure pain interference. *Pain* 2010 Jul;150(1):173-182 [FREE Full text] [doi: [10.1016/j.pain.2010.04.025](https://doi.org/10.1016/j.pain.2010.04.025)] [Medline: [20554116](https://pubmed.ncbi.nlm.nih.gov/20554116/)]
45. Jensen RE, King-Kallimani BL, Sexton E, Reeve BB, Moinpour CM, Potosky AL, et al. Measurement properties of PROMIS sleep disturbance short forms in a large, ethnically diverse cancer cohort. *Psychol Test Assess Model* 2016;58(2):353 [FREE Full text]
46. Yu L, Buysse DJ, Germain A, Moul DE, Stover A, Dodds NE, et al. Development of short forms from the PROMIS™ sleep disturbance and sleep-related impairment item banks. *Behav Sleep Med* 2011 Dec 28;10(1):6-24 [FREE Full text] [doi: [10.1080/15402002.2012.636266](https://doi.org/10.1080/15402002.2012.636266)] [Medline: [22250775](https://pubmed.ncbi.nlm.nih.gov/22250775/)]
47. Holmes TH, Rahe RH. The social readjustment rating scale. *J Psychosom Res* 1967 Aug;11(2):213-218. [doi: [10.1016/0022-3999\(67\)90010-4](https://doi.org/10.1016/0022-3999(67)90010-4)]
48. Patel AT, Duncan PW, Lai SM, Studenski S. The relation between impairments and functional outcomes poststroke. *Arch Phys Med Rehabil* 2000 Oct;81(10):1357-1363. [doi: [10.1053/apmr.2000.9397](https://doi.org/10.1053/apmr.2000.9397)] [Medline: [11030501](https://pubmed.ncbi.nlm.nih.gov/11030501/)]
49. Jørgensen HS, Nakayama H, Raaschou HO, Olsen TS. Recovery of walking function in stroke patients: the Copenhagen stroke study. *Arch Phys Med Rehabil* 1995 Jan;76(1):27-32. [doi: [10.1016/S0003-9993\(95\)80038-7](https://doi.org/10.1016/S0003-9993(95)80038-7)] [Medline: [7811170](https://pubmed.ncbi.nlm.nih.gov/7811170/)]
50. Feigin VL, Roth GA, Naghavi M, Parmar P, Krishnamurthi R, Chugh S, Global Burden of Diseases, Injuries and Risk Factors Study 2013 and Stroke Experts Writing Group. Global burden of stroke and risk factors in 188 countries, during

- 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet Neurol* 2016 Aug;15(9):913-924. [doi: [10.1016/S1474-4422\(16\)30073-4](https://doi.org/10.1016/S1474-4422(16)30073-4)] [Medline: [27291521](https://pubmed.ncbi.nlm.nih.gov/27291521/)]
51. Myint PK, Luben RN, Wareham NJ, Bingham SA, Khaw KT. Combined effect of health behaviours and risk of first ever stroke in 20,040 men and women over 11 years' follow-up in Norfolk cohort of European prospective investigation of cancer (EPIC Norfolk): prospective population study. *Br Med J* 2009 Feb 19;338:b349 [FREE Full text] [doi: [10.1136/bmj.b349](https://doi.org/10.1136/bmj.b349)] [Medline: [19228771](https://pubmed.ncbi.nlm.nih.gov/19228771/)]
 52. Global Burden of Metabolic Risk Factors for Chronic Diseases Collaboration (BMI Mediated Effects), Lu Y, Hajifathalian K, Ezzati M, Woodward M, Rimm EB, et al. Metabolic mediators of the effects of body-mass index, overweight, and obesity on coronary heart disease and stroke: a pooled analysis of 97 prospective cohorts with 1.8 million participants. *Lancet* 2014 Mar 15;383(9921):970-983 [FREE Full text] [doi: [10.1016/S0140-6736\(13\)61836-X](https://doi.org/10.1016/S0140-6736(13)61836-X)] [Medline: [24269108](https://pubmed.ncbi.nlm.nih.gov/24269108/)]
 53. Cleveland S, Driver S, Swank C, Macklin S. Classifying physical activity research following stroke using the behavioral epidemiologic framework. *Top Stroke Rehabil* 2015 Aug;22(4):289-298. [doi: [10.1179/1074935714Z.0000000043](https://doi.org/10.1179/1074935714Z.0000000043)] [Medline: [26258454](https://pubmed.ncbi.nlm.nih.gov/26258454/)]
 54. Green T, Haley E, Eliasziw M, Hoyte K. Education in stroke prevention: efficacy of an educational counselling intervention to increase knowledge in stroke survivors. *Can J Neurosci Nurs* 2007;29(2):13-20. [Medline: [18240627](https://pubmed.ncbi.nlm.nih.gov/18240627/)]
 55. Kono Y, Yamada S, Yamaguchi J, Hagiwara Y, Iritani N, Ishida S, et al. Secondary prevention of new vascular events with lifestyle intervention in patients with noncardioembolic mild ischemic stroke: a single-center randomized controlled trial. *Cerebrovasc Dis* 2013;36(2):88-97. [doi: [10.1159/000352052](https://doi.org/10.1159/000352052)] [Medline: [24029303](https://pubmed.ncbi.nlm.nih.gov/24029303/)]
 56. White JH, Bynon BL, Marquez J, Sweetapple A, Pollack M. 'Masterstroke: a pilot group stroke prevention program for community dwelling stroke survivors'. *Disabil Rehabil* 2013 Jun;35(11):931-938. [doi: [10.3109/09638288.2012.717578](https://doi.org/10.3109/09638288.2012.717578)] [Medline: [23641954](https://pubmed.ncbi.nlm.nih.gov/23641954/)]
 57. Betts AC, Froehlich-Grobe K. Accessible weight loss: adapting a lifestyle intervention for adults with impaired mobility. *Disabil Health J* 2017 Jan;10(1):139-144. [doi: [10.1016/j.dhjo.2016.06.004](https://doi.org/10.1016/j.dhjo.2016.06.004)] [Medline: [27431768](https://pubmed.ncbi.nlm.nih.gov/27431768/)]
 58. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007 Aug;39(8):1423-1434. [doi: [10.1249/mss.0b013e3180616b27](https://doi.org/10.1249/mss.0b013e3180616b27)] [Medline: [17762377](https://pubmed.ncbi.nlm.nih.gov/17762377/)]
 59. Sallis RE. Exercise is medicine and physicians need to prescribe it!. *Br J Sports Med* 2009 Jan;43(1):3-4. [doi: [10.1136/bjism.2008.054825](https://doi.org/10.1136/bjism.2008.054825)] [Medline: [18971243](https://pubmed.ncbi.nlm.nih.gov/18971243/)]
 60. Gickman D, Parker L, Sim LJ, del Valle Cook H, Miller EA, Committee on Accelerating Progress in Obesity Prevention, Food and Nutrition Board, Institute of Medicine. *Accelerating Progress in Obesity Prevention: Solving the Weight of the Nation*. Washington, DC: National Academies Press; 2012.
 61. Driver S, Ede A, Dodd Z, Stevens L, Warren AM. What barriers to physical activity do individuals with a recent brain injury face? *Disabil Health J* 2012 Apr;5(2):117-125. [doi: [10.1016/j.dhjo.2011.11.002](https://doi.org/10.1016/j.dhjo.2011.11.002)] [Medline: [22429546](https://pubmed.ncbi.nlm.nih.gov/22429546/)]
 62. Damush TM, Plue L, Bakas T, Schmid A, Williams LS. Barriers and facilitators to exercise among stroke survivors. *Rehabil Nurs* 2007;32(6):253-60,262. [doi: [10.1002/j.2048-7940.2007.tb00183.x](https://doi.org/10.1002/j.2048-7940.2007.tb00183.x)] [Medline: [18065147](https://pubmed.ncbi.nlm.nih.gov/18065147/)]
 63. Nicholson SL, Donaghy M, Johnston M, Sniehotta FF, van Wijck F, Johnston D, et al. A qualitative theory guided analysis of stroke survivors' perceived barriers and facilitators to physical activity. *Disabil Rehabil* 2014;36(22):1857-1868. [doi: [10.3109/09638288.2013.874506](https://doi.org/10.3109/09638288.2013.874506)] [Medline: [24377329](https://pubmed.ncbi.nlm.nih.gov/24377329/)]
 64. Simpson LA, Eng JJ, Tawashy AE. Exercise perceptions among people with stroke: barriers and facilitators to participation. *Int J Ther Rehabil* 2011 Sep 6;18(9):520-530 [FREE Full text] [doi: [10.12968/ijtr.2011.18.9.520](https://doi.org/10.12968/ijtr.2011.18.9.520)] [Medline: [23255881](https://pubmed.ncbi.nlm.nih.gov/23255881/)]
 65. Hinnant L, Razi S, Lewis R, Sun A, Alva M, Hoerger T, et al. Evaluation of the health care innovation awards: Community resource planning, prevention and monitoring, annual report. North Carolina: RTI; 2015.

Abbreviations

ACL: Administration for Community Living

AHA: American Heart Association

BMI: body mass index

CDC: Centers for Disease Control and Prevention

CDC-DPRP: Centers for Disease Control and Prevention-National Diabetes Prevention Recognition Program

CVA: cerebrovascular accident

GLB: Group Lifestyle Balance

HHS: Health and Human Services

NIDILRR: National Institute on Disability, Independent Living, and Rehabilitation Research

RCT: randomized controlled trial

Edited by G Eysenbach; submitted 10.04.19; peer-reviewed by J Bishop, K Best; comments to author 27.06.19; revised version received 11.07.19; accepted 19.07.19; published 18.10.19

Please cite as:

Driver S, Swank C, Froehlich-Grobe K, McShan E, Calhoun S, Bennett M

Weight Loss After Stroke Through an Intensive Lifestyle Intervention (Group Lifestyle Balance-Cerebrovascular Accident): Protocol for a Randomized Controlled Trial

JMIR Res Protoc 2019;8(10):e14338

URL: <https://www.researchprotocols.org/2019/10/e14338>

doi: [10.2196/14338](https://doi.org/10.2196/14338)

PMID: [31628790](https://pubmed.ncbi.nlm.nih.gov/31628790/)

©Simon Driver, Chad Swank, Katherine Froehlich-Grobe, Evan McShan, Stephanie Calhoun, Monica Bennett. Originally published in JMIR Research Protocols (<http://www.researchprotocols.org>), 18.10.2019. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Research Protocols, is properly cited. The complete bibliographic information, a link to the original publication on <http://www.researchprotocols.org>, as well as this copyright and license information must be included.