

Protocol

Rationale and Design of an Ecological Momentary Assessment Study Examining Predictors of Binge Eating Among Sexual Minority and Heterosexual Young Women: Protocol for the Health and Experiences in Real Life (HER Life) Study

Kristin E Heron^{1,2}, PhD; Abby L Braitman^{1,2}, PhD; Charlotte A Dawson², MS; Cassidy M Sandoval², MA; Lauren V Butler¹, BS; Alicia Moulder¹, BS; Robin J Lewis^{1,2}, PhD

¹Department of Psychology, Old Dominion University, Norfolk, VA, United States

²Virginia Consortium Program in Clinical Psychology, Norfolk, VA, United States

Corresponding Author:

Kristin E Heron, PhD
Department of Psychology
Old Dominion University
250 Mills Godwin Building
Norfolk, VA, 23529
United States
Phone: 1 757 683 5214
Email: kheron@odu.edu

Abstract

Background: Previous research has identified health disparities between sexual minority and heterosexual women, including increased rates of obesity and binge eating in sexual minority women. Established predictors of binge eating behavior include negative emotions and sociocultural processes; however, these studies are generally conducted in samples of young women where sexual identity is not known or reported. There is a dearth of research evaluating how sexual minority-specific factors (eg, minority stress and connectedness to the lesbian, gay, bisexual, transgender, and queer community) may affect binge eating in sexual minority women. In addition, no studies have examined these processes in racially diverse samples or considered how intersecting minority identities (eg, Black and sexual minority) may affect eating behaviors.

Objective: The Health and Experiences in Real Life (HER Life) Project aims to clarify real-world predictors of binge eating in young heterosexual and sexual minority women using ecological momentary assessment. The role of affective, social, and health behavior factors in binge eating will be examined for all women (aim 1), and sexual minority-specific predictors will also be considered for sexual minority women participants (aim 2). Person-level moderators of race, body- and eating-related factors, and sexual minority-specific factors will also be examined to better understand how real-world binge eating predictors may differ for various demographic groups (aim 3).

Methods: Researchers aim to recruit 150 sexual minority and 150 heterosexual women from across the United States, including at least 50 Black women for each group, using web-based recruitment methods. The eligibility criteria include identifying as a woman, being aged between 18 and 30 years, and having had at least two binge eating episodes in the last 2 weeks. Participants must endorse being only or mostly attracted to men (considered heterosexual) or only or mostly attracted to women or having a current or most recent female partner (considered sexual minority). Eligible participants complete an initial web-based baseline survey and then 14 days of ecological momentary assessment involving the completion of a morning and before-bed survey and 5 prompted surveys per day as well as a user-initiated survey after binge eating episodes. The data will be analyzed using a series of multilevel models.

Results: Data collection started in February 2021. We have currently enrolled 129 sexual minority women and 146 heterosexual women. Data collection is expected to conclude in fall 2022.

Conclusions: The Health and Experiences in Real Life Project aims to elucidate potential differences between sexual minority and heterosexual women in within-person factors predicting binge eating and inform eating disorder interventions for sexual minority women. The challenges in recruiting sexual minority women, including the determination of eligibility criteria and considerations for remote data collection, are discussed.

International Registered Report Identifier (IRRID): DERR1-10.2196/41199

(*JMIR Res Protoc* 2022;11(10):e41199) doi: [10.2196/41199](https://doi.org/10.2196/41199)

KEYWORDS

sexual minority women; ecological momentary assessment; binge eating; sexual minority stress; negative affective states; mobile phone

Introduction

Background

Obesity and obesity-related conditions are the leading causes of death in the United States [1], making them a significant public health concern. On the basis of several recent national reports [2,3], there are disparities in the health and well-being of sexual minority individuals (eg, gay, lesbian, and bisexual) as compared with their heterosexual peers in a range of health conditions and health behaviors. In particular, large-scale studies suggest that sexual minority women are at least twice as likely to be obese as heterosexual women [4,5], and studies of young women have similarly found higher rates of being overweight or obese in young lesbian women (35.2%) than in young heterosexual women (22.8%) [6]. There is also evidence that rates of binge eating, defined as overeating with loss of control of eating, are similarly higher in sexual minority individuals. For example, in a study of nearly 14,000 people, young adult lesbian women were twice as likely to binge eat as heterosexual women [7]. Despite these documented disparities, relatively little is known about the contributing factors, which is a critical first step in developing interventions to address these health concerns among sexual minority women.

In working toward increased understanding of factors that contribute to higher obesity and binge eating rates in sexual minority women, the goal of this study is to use ecological momentary assessment (EMA) to identify general and minority-specific real-world predictors of binge eating. EMA involves the use of repeated assessments of people's current behaviors, states, and experiences as they are going about their lives [8]. EMA approaches provide unique advantages over other research methods (eg, cross-sectional and traditional longitudinal methods), including greater real-world generalizability and reduced concerns regarding memory biases associated with retrospective recall that are generally present in traditional measures. In addition, as EMA uses repeated assessments (typically multiple times throughout the day), researchers can examine dynamic within-person processes and answer research questions that cannot be addressed using other methods [8,9]. For example, most previous research on binge eating among sexual minority women has used cross-sectional designs, which can address questions regarding between-person processes (eg, do women who are more depressed also binge eat more?). EMA approaches provide an opportunity to answer questions regarding *within-person* processes (eg, when women experience a more depressed mood, are they more likely to binge eat?). This study—called the Health and Experiences in Real Life (HER Life) Project—will examine within-person associations between a range of general predictors of binge eating (ie, affect, body dissatisfaction, social processes, and

health behaviors) in young heterosexual and sexual minority women as well as sexual minority-specific factors (eg, minority stress and discrimination) among young sexual minority women. The existing empirical evidence and theoretical framework supporting this study are described in the following sections.

General Factors Associated With Binge Eating in Daily Life (Aim 1)

The factors influencing disordered eating are complex, and researchers have highlighted the potential utility of EMA to study disordered eating, including binge eating, in natural settings [10]. EMA research shows that disordered eating, including binge eating, is associated with affective states (eg, negative mood, stress, and body dissatisfaction) [11,12], social experiences (eg, interactions and media use), and health behaviors (eg, alcohol use and physical activity) in studies of young women where sexual orientation was not assessed or reported (hereafter referred to as general samples). However, with few exceptions [13], EMA studies have yet to consider how affective states, social processes, and health behaviors in daily life are associated with sexual minority women's binge eating, which is a primary aim of this study.

The role of negative emotions (eg, negative affect and stress) in binge eating is often examined as a way of evaluating the affect regulation model of eating disorders [14]. A meta-analysis of EMA studies evaluating the affect regulation model of binge eating found that negative emotions increased leading up to a binge eating episode, with more mixed findings regarding negative affect after binge eating episodes [12]. Negative feelings about one's body, or body dissatisfaction, have also been examined via EMA in general samples of young women but, with few exceptions [15], studies have generally focused on associations between body dissatisfaction and social processes, not disordered eating behaviors [16-18]. Although these studies show that negative mood and body dissatisfaction may be associated with binge eating in daily life, this work is largely based on general samples of young women, and whether and how these processes operate in sexual minority women is less clear. There is some preliminary evidence demonstrating that negative affect is associated with binge eating at the day level in a daily diary study of 30 lesbian women [13], but there are no EMA studies of body dissatisfaction in sexual minority women. Cross-sectional studies designed to answer between-person research questions have produced mixed findings, with some suggesting that body dissatisfaction is similar in sexual minority and heterosexual women [19] and others finding that larger body sizes are more acceptable among sexual minority social groups and, thus, body dissatisfaction may be lower [5]. However, as described previously, this study will fill gaps in the literature by assessing negative affect, stress, and body dissatisfaction in daily life to examine how they are

associated with binge eating among both young sexual minority and heterosexual women.

There is a long history of research demonstrating the contribution of sociocultural processes to disordered eating behaviors [20,21]. In fact, the sociocultural model of disordered eating suggests that there are social pressures to be thin (and, increasingly, to be fit or toned) in Western countries, which are communicated directly and indirectly through the media, peers, and family and can become internalized (ie, thin ideal internalization) [22]. In recent years, several EMA studies have examined how social experiences are associated with body dissatisfaction and eating behaviors in daily life. For example, in a general sample of 121 women aged 18 to 40 years, women reported greater body dissatisfaction in social situations (ie, when they had had an interaction in the previous 30 minutes) than when alone and particularly when they perceived that the interaction quality was lower [18]. To our knowledge, studies on the effect of social processes in daily life on eating behaviors have been conducted only among general samples of young women and, thus, it remains unclear whether sexual minority women experience the same sociocultural influences as heterosexual women in daily life. Cross-sectional research suggests that larger body sizes may be more acceptable in sexual minority women's social groups and relationships [5,23] and, thus, they may be less susceptible to thin ideals that they are exposed to in social interactions and the media. However, as described by the dual identity framework, there may also be commonalities between sexual minority and heterosexual women [24]. According to this framework, sexual minority women are influenced, as women, by the mainstream (heterosexual) community and, as sexual minority individuals, by the sexual minority community. Thus, sexual minority women's eating behaviors will likely be influenced by general social processes but potentially to a lesser extent than heterosexual women. The HER Life Project will explore social influences on both sexual minority and heterosexual women in daily life, thus allowing for a better understanding of how these dual influences may operate for young sexual minority women.

In addition to the demonstrated importance of sociocultural considerations for understanding disordered eating, cross-sectional and meta-analytic studies have shown that hazardous alcohol use is associated with disordered eating in general samples of young adults [25] and among lesbian women. For instance, a cross-sectional survey found that lesbian women with obesity were more likely to report heavy drinking compared with lesbian women who were not overweight or obese [26]. Research on the associations between binge eating and physical activity are more equivocal [27], but 1 study with a general sample of young women found that those with binge eating disorder reported less physical activity compared with weight-matched women who did not have binge eating disorder [28]. Even less is known about physical activity and binge eating in sexual minority women, but 1 study of lesbian and bisexual women found that being overweight or obese was associated with less frequent exercise [29]. It is clear that there is a dearth of research on the associations among alcohol use, physical activity, and binge eating, especially in sexual minority women. This is particularly notable because of the documented

disparities in alcohol use and physical activity, with sexual minority women engaging in more hazardous drinking [30-32] and less physical activity [33,34] than heterosexual women. This study will assess alcohol use and physical activity in daily life, which will allow for a real-world examination of how these health behaviors may operate among sexual minority and heterosexual young women, and explore potential associations with binge eating.

Sexual Minority–Specific Factors Associated With Binge Eating in Daily Life (Aim 2)

In addition to the aforementioned general factors that may influence binge eating, sexual minority women face unique experiences and stressors because of their marginalized and stigmatized status in society. This phenomenon has been described in minority stress theories by Meyer [35] and Hatzenbuehler [36]. These theories complement each other in describing how general and minority-specific stressors can affect mental health and well-being. For example, the minority stress theory by Meyer argues that sexual minority individuals experience stressors related to their minority status. These stressors include chronic and acute external events (discrimination, harassment, invalidations, and vicarious experiences through other sexual minority individuals) and internal stressors (concealment of sexual identity, expected rejection, and internalization of negative societal messages about sexual minorities [internalized heterosexism]) that can negatively affect mental health. The psychological mediation framework by Hatzenbuehler extended these ideas to suggest that general (eg, coping, emotion regulation, and social support) and minority-specific (eg, distal and proximal minority stressors) experiences operate together to influence sexual minority individuals' well-being. These theories have been extended from their initial focus on understanding mental health to also consider the impact of minority stressors on physical health and health behaviors.

In recent years, minority stress theories have been used to identify potential mechanisms to explain health disparities between sexual minority and heterosexual individuals, including disordered eating. Cross-sectional studies of sexual minority women have found between-person associations between heterosexual experiences (eg, harassment, rejection, and sexual orientation discrimination) and disordered eating [37-39]. More recently, there have been several daily diary and EMA studies that have assessed within-person associations between minority stressors and disordered eating behaviors at the day or moment level. For example, in a sample of 30 young lesbian women who reported binge eating, researchers found that, on days when the women reported more discrimination, they also reported more binge eating (and this effect occurred in part through negative affect) [13]. In an EMA study of 55 sexual minority women who were overweight or obese, Panza et al [40] similarly found that those with higher baseline levels of internalized homophobia and sexual orientation concealment reported more binge eating and overeating during a 5-day EMA period. Taken together, sexual minority stress theories and empirical research suggest that sexual minority stress is associated with poor mental health, with substantially fewer studies considering the effects on binge eating. The goal of this study is to examine the

within-person direct and indirect (through negative affect) effects of sexual minority stressors on binge eating in a subsample of sexual minority women. In other words, this study allows us to consider whether, at times when sexual minority women experience a minority stressor, that experience is either concurrently or subsequently associated with binge eating in daily life. Such information may improve the understanding of the potential mechanisms whereby sexual minority women experience more disordered eating behaviors as compared with heterosexual women.

Risk and Protective Factors for Binge Eating (Aim 3)

Given that theories on binge eating and sexual minority health and health disparities are complex, with factors influencing these behaviors at various *levels*, an additional aim of this study is to consider how *between-person* factors may strengthen or weaken the aforementioned *within-person* associations. Three broad groups of moderators will be considered in this study: race, body- and eating-related factors, and sexual minority-specific factors. Considering these cross-level moderators can be useful for clarifying previous conflicting findings and provide a more complete picture of both *when* and *for whom* various real-world binge eating predictors operate.

Although cross-sectional research shows that sexual minority women are more likely to report binge eating compared with heterosexual women [7,41,42], many of these studies were conducted in samples that were not racially diverse. There is some conflicting research regarding potential racial differences in binge eating between Black and White women in the United States. Some studies in general samples of young women have found no differences in disordered eating (including binge eating) between Black and White women [43,44] but, in a general sample of women with binge eating disorder, Black women reported 1.5 times as many binge eating episodes per week compared with White women [45]. However, taken together, these studies fail to consider how multiple minority identities (ie, being a Black sexual minority woman) might interact to exacerbate health behavior risks, and there is scant research examining the effect of the intersection of sexual and racial identities on binge eating. In fact, we found only 1 study that assessed binge eating in a small, racially diverse sample of lesbian and bisexual women. This cross-sectional study found that 1.6% of White heterosexual women (n=63) reported binge eating at least twice per week compared with rates of 4.5% for White lesbian and bisexual women (n=67) and 3.1% for Black lesbian and bisexual women (n=64) [46]. Importantly, this study was cross-sectional and had a small sample. Therefore, to extend these past findings, in the HER Life Project, we powered to test for Black-White race moderator effects of the within-person associations between the general and minority-specific predictors of binge eating outlined in the first 2 study aims, allowing us to examine how intersecting minority statuses (ie, being Black and of a sexual minority) may uniquely shape eating behavior in daily life.

In addition to considering how women's racial identity may moderate within-person associations, this study will consider 4 body- and eating-related variables: BMI, person-level (baseline) body dissatisfaction, disordered eating, and thin norm

internalization. These variables were selected based on past research demonstrating that individuals higher or lower in these constructs may exhibit different associations between our predictors of interest and binge eating [17].

In addition, this study considers sexual minority-specific moderators of EMA associations, which will be considered among the subsample of sexual minority women. In particular, we will evaluate 3 protective factors (positive sexual minority identity; lesbian, gay, bisexual, transgender, and queer [LGBTQ+] community connection; and identification with the LGBTQ+ community) and one risk factor (lifetime sexual minority discrimination). Previous cross-sectional research suggests that connection to and identification with the sexual minority community may buffer against the deleterious effects of sexual minority stress [47], including perceived stigma and depressive symptoms [48]. Furthermore, in a cross-sectional study of lesbian, gay, and bisexual young adults, social support (eg, from the sexual minority community) attenuated the associations between minority stress and emotional distress [49]. Conversely, lifetime discrimination may be a risk factor for maladaptive health behaviors. In a daily diary study of Black college students, perceived lifetime racial discrimination moderated the associations between negative mood and alcohol use [50], providing evidence that (person-level) lifetime racial discrimination influences daily health behaviors. Taken together, the third aim of this study is to consider a variety of person-level moderators of the within-person associations explored in the first 2 study aims.

Overview of Study Aims

The broad objective of the HER Life Project is to enhance the understanding of the real-world predictors of binge eating among young sexual minority and heterosexual women. To address this objective, this study is organized around 3 aims.

Aim 1: To Examine the Role of Affective, Social, and Health Behavior Factors in Binge Eating

We hypothesize that, when sexual minority and heterosexual women experience higher levels of negative emotion variables (negative affect, general stress, and body dissatisfaction), they engage in more binge eating (hypothesis 1a). We also expect that social experiences (eg, social interactions and media use) will be associated with more binge eating for both groups (hypothesis 1b), but the associations will be stronger for heterosexual women than for sexual minority women (hypothesis 1c). A priori hypotheses for binge eating, alcohol use, and physical activity associations are not made because of limited previous EMA research.

Aim 2: To Examine the Associations Between Sexual Minority Stress and Binge Eating Among Sexual Minority Women

Direct within-person associations are hypothesized such that, when sexual minority women experience more sexual minority stress, they will engage in more binge eating (hypothesis 2a). Consistent with theory, we also expect that negative affect will mediate the association between sexual minority stress and binge eating (hypothesis 2b). These aims will be tested within the subsample of sexual minority women recruited for this study.

Aim 3: To Explore Race and Other Person-Level Moderators of Associations in Aims 1 and 2

Race, body- and eating-related factors (BMI, body dissatisfaction, disordered eating, and thin norm internalization), and sexual minority-specific factors (lifetime discrimination, positive sexual minority identity, and community identification and connection) will be explored as moderators of EMA associations in aims 1 and 2. Given the scant previous research, no a priori hypotheses are specified, but the study is powered to test for Black-White group differences for aims 1 and 2 to explore the intersection of multiple minority identities.

Methods

Project Overview

The HER Life Project is an ongoing EMA study of young sexual minority and heterosexual women who binge eat. Young women aged 18 to 30 years are the focus of this study given that they report high levels of binge eating [4,5] and, thus, represent an at-risk group. We selected women aged 18 to 30 years as, although “young adulthood” has no clear definition, there is evidence that sexual identity continues to develop during this period, and data suggest associations among sexual identity, negative affect, and disordered eating up to the age of approximately 30 years [51]. Participants are being recruited from across the United States, and data collection is occurring entirely remotely as we are using web-based surveys and a smartphone app to deploy EMA surveys. Participants receive and review the written and video study procedure descriptions before beginning the study. They complete a web-based screening and baseline survey before beginning the 2-week EMA data collection period and conclude their participation with an end-of-study survey. During the EMA period, participants complete daily morning and before-bed surveys, 5 prompted surveys each day, and a user-initiated binge eating survey after binge eating episodes.

Ethics Approval

The Old Dominion University Institutional Review Board approved all the study procedures (project 1362990).

Power Analysis

A 3-step power analysis was conducted to determine sample size. First, a regression power analysis using G*Power (version 3.1.9) [52] was conducted, powering the hardest-to-detect effects (aim 1 hypothesis 1c: sexual identity moderation; aim 3: race moderation) and the examinations using only sample subsets (aim 2: direct effects, sexual minority women only). To achieve a power of 0.80 using an α of .05, 227 independent observations are required for a small to medium effect ($f^2=0.035$) for moderation examinations with 5 predictors (to allow for relevant covariates). Specifying the same power level, α , and effect size with 3 predictors for aim 2 main effects (to allow for relevant covariates), 227 independent observations among sexual minority women only would be necessary. Second, we applied a formula to account for the multiple correlated observations for each person and identify the number of actual participants needed [53]. This equation accounts for level-1 (assessments or moments) and level-2 (participants) sample sizes and the

intraclass correlation coefficient (ICC; within-person relatedness). On the basis of past EMA studies of similar constructs among young women [54,55], a conservative estimate of the average number of assessments per person (n_{L1}) is 25, reflecting constructs only asked once or twice per day (as opposed to all prompts), approximately 70% compliance, and the reporting of approximately 6 binge eating episodes. If 100 sexual minority and 100 heterosexual women are recruited, this would result in approximately 5000 total observations (2500 from sexual minority women). We expect ICC values from 0.33 to 0.43 for binge eating. Using the West et al [53] formula and past studies, n_{L2} of 100 sexual minority and 100 heterosexual women would result in an N-effective of 280 within sexual identity if the ICC is 0.33 and an N-effective of 221 within sexual identity if the ICC is 0.43. This means that, although there are 2500 assessments from sexual minority women, because of the degree of relatedness of observations within each individual, these are equivalent to 221 independent pieces of information. Thus, recruiting 200 women should provide sufficient power for these aims. In addition, to detect small to medium mediation effects (aim 2) using empirical bias-corrected bootstrap CIs, 148 independent observations should be sufficient [56]. Finally, to power for aim 3, half of the 100 participants in each sexual identity group ($n=50$) will need to identify as Black to optimize power. However, given the demographic representation in previous work examining eating behaviors in young women using EMA [54,55], it is expected that we will need to enroll >100 women of each identity to have 50 Black women in each group even if deliberately oversampling. To ensure that we are sufficiently powered to test all aims and have similar numbers in both groups, we aim to enroll 150 sexual minority women (50 Black) and 150 heterosexual women (50 Black).

Participant Selection

Participants are being recruited using various web-based sources, including from web-based research panels, social media, and lists of people who participated in past studies and expressed an interest in future research opportunities. We are working with several market research firms who have access to web-based panels of people who are interested in participating in web-based research. Community Marketing & Insights is an LGBTQ+ market research firm that manages a proprietary panel of individuals who identify within the LGBTQ+ community and are interested in participating in web-based studies. Marketing Systems Group and Qualtrics International Inc are also assisting with accessing existing panels of individuals who regularly participate in web-based studies, primarily to recruit heterosexual women. We have also placed advertisements on social media (eg, Facebook and Instagram) using the Meta advertising platform and through advertisements on our laboratory social media page.

To be eligible to take part in the study, participants must (1) identify as women (from a multi-select question, selecting woman or woman *and* some combination of genderqueer, nonconforming, or nonbinary and also indicating that they were assigned female at birth), (2) be aged between 18 and 30 years (inclusive), (3) report binge eating (ie, overeating with a feeling

of loss of control of eating) at least two times in the past 2 weeks, (4) not be currently receiving treatment for an eating disorder, and (5) have a schedule that allows for answering surveys during daytime hours. In addition, the participants must also meet either the sexual minority or heterosexual criteria. Women are included in the sexual minority sample if they report that they are (1) only or mostly attracted to women or (2) equally attracted to men and women or attracted to people regardless of their gender identity *and* also have a current or most recent romantic partner who identifies as a woman. Women are included in the heterosexual sample if they report that they are only or mostly attracted to men.

Study Procedures

Recruitment

For participants being recruited through the marketing research firms, the firm makes initial contact with potential participants and administers an initial web-based screening survey. Contact information for the eligible participants is then provided to the research team. At this time, participants are contacted via email and screened directly by the research team using a brief web-based screening survey. For participants being recruited through social media and lists of past research participants interested in research studies, potential participants are provided directly with a web-based screening survey and screened by our team.

Informed Consent Process and Baseline Survey

Eligible participants are provided with additional information about the study during the informed consent process. First, a 3-minute video providing a general overview of the study's purpose, procedures, and participation requirements (eg, downloading a smartphone app) and compensation is emailed to potentially eligible participants. Second, participants are provided with information about how to download the smartphone app needed to complete the EMA surveys. Participants confirm their interest in participating by downloading the smartphone app and notifying the research team that they have done so. The LifeData (LifeData, LLC) survey software and RealLife Exp (LifeData, LLC) app are being used for this study. This software allows for custom EMA surveys and alarm schedules to be created, and the surveys can be deployed on any Android or Apple smartphone using the RealLife Exp app. Plans are in place to provide a phone to participants if they do not have an Android or Apple phone that they are willing or able to use for the study (none have requested this option to date). Participants who do not respond to these emails are sent up to 3 reminders to review the video and download the smartphone app. Additional details regarding the email reminder system used throughout the study process are provided in the following sections. Once participants confirm their interest in proceeding with the study (as evidenced by downloading the app to their phone), they are sent a web-based link to complete an electronic informed consent form, which is immediately followed by the baseline survey, which takes approximately 45 minutes to complete.

EMA Surveys

After completing the baseline survey, participants receive an email from our team with instructions on how to download the study-specific EMA smartphone surveys (within the app they previously downloaded) and a document with frequently asked questions and answers about study procedures. Once participants download the study-specific surveys, they complete an initial "startup" session where they enter their study-specific identification number and watch 2 additional brief training videos created by the research team. The first video provides additional information about when and how to complete each type of EMA survey and is approximately 3 minutes long. The second video provides a definition and examples of a binge eating episode to help participants in identifying their binge eating and improve understanding of when to initiate a binge eating survey; this video is approximately 2 minutes long. These informational videos and the document with frequently asked questions and answers are accessible to participants within the smartphone app throughout the entirety of the EMA portion of the study.

The day after the study-specific EMA surveys are downloaded, participants begin to receive 5 prompted surveys per day for the next 14 days at semirandom times between 9 AM and 9 PM each day. In addition to the prompted surveys, participants complete 3 different user-initiated surveys: an after binge eating survey, a morning survey, and a before-bed survey. They are instructed to complete the post-binge eating survey after a binge eating episode; as described previously, participants receive written and video instructions defining, describing, and providing examples of a binge eating episode. They are instructed to self-initiate the morning survey immediately after waking up and the before-bed survey just before going to sleep at night, but the app also provides a reminder notification for each (at 10 AM for the morning survey and 9 PM for the before-bed survey).

End-of-Study Survey

After day 14 of the EMA surveys, participants receive an email including a link to complete a survey regarding their experience as research participants in this study. Additional information on this measure can be found in the Measures section.

Compensation

Participants can earn up to US \$150 for taking part in this study. They receive US \$20 for completing the baseline survey and US \$10 for completing the end-of-study survey. Each week of the EMA, they receive US \$40 (a total of US \$80). To incentivize compliance, they can also receive a US \$20 bonus each week if they complete 80% of the daily surveys (averaging a morning survey, 4 prompted surveys, and a before-bed survey per day). Within a week of completing the end-of-study survey, participants receive an email thanking them for taking part and notifying them of their compensation amount. Participants receive an electronic gift card (selected from several options) via email at the conclusion of their participation.

Email Reminder and Check-in Systems

Given that this study is being conducted entirely remotely and participants need to move through various steps to enroll in and

complete the study, we developed a system for reminding participants who do not respond to initial emails or complete tasks as expected. Throughout each stage of the process—screening, completing the baseline survey, beginning the EMA surveys, and completing the end-of-study survey—project staff send participants up to 2 reminders, approximately 2 to 3 days apart, to encourage participation. If participants do not respond to these reminders, a final email is sent by the study principal investigator (PI; KH) to inquire about questions and encourage their continued participation.

In addition to monitoring the overall flow throughout the study, we carefully track compliance with the EMA surveys. Participants' completion of the daily surveys is tracked and recorded each day using a daily tracking spreadsheet. Participants are sent a general check-in email on day 2 or 3, day 7 or 8, and day 13 or 14 of the EMA portion of the study to thank participants for their completion thus far, offer the opportunity to ask questions, and encourage continued compliance. Participants demonstrating a pattern of noncompliance (eg, no prompted surveys completed in a day, no morning or before-bed surveys completed in a day, or many prompted surveys not completed) receive a noncompliance email. Noncompliance emails are sent to participants to advise them as to which surveys they are not completing on a regular basis, check whether they are experiencing difficulty completing the daily surveys, and encourage participation.

Measures

The measures in the baseline survey are described in [Table 1](#) and are grouped together by construct. Participants generally complete the same baseline survey, with 2 exceptions. First, there are several specific questionnaires for sexual minority women regarding minority stressors (eg, discrimination and rejection), identity, and LGBT community connection. To balance survey length and content, heterosexual women complete questionnaires regarding their identity as women and their general social support. Second, as we are interested in assessing alcohol use but did not require participants to drink alcohol to participate in this study, we ask about alcohol use only of those women who drank in the past 30 days and administer a series of questionnaires regarding reasons for not drinking and general coping to nondrinkers to similarly balance for time.

The measures in the 4 different EMA surveys (morning, before bed, after binge eating, and prompted) are described in [Table 2](#). Items were either drawn directly from existing EMA scales or past studies or were adapted from non-EMA measures and developed for the purposes of this study in instances where no EMA measure of a construct existed.

The end-of-study survey collects information about participants' experiences in the study, interest in and willingness to be involved in future mobile health interventions regarding physical and mental health, and COVID-19–related questions. [Table 3](#) provides a description of the constructs included in the end-of-study survey.

Table 1. Baseline measures.

Construct and description	Measure name
Demographics—age, sexual orientation (identity, attraction, and behavior), height, weight, address, geographic location, relationship status and length, employment status, education level, average individual income, and finances	N/A ^a
Sexual identity—self-identity, identity disclosure, and “coming out” (for sexual minority women only)	N/A
Eating and body image	
Eating pathology (body dissatisfaction, binge eating, cognitive restraint, purging, restricting, and excessive exercise)	Eating Pathology Symptoms Inventory [57]
Diagnostic criteria for eating disorder behaviors	Eating Disorder Examination Questionnaire 6.0—selected items [58]
Trait-level body dissatisfaction	Body Shape Questionnaire [59]
Societal and interpersonal norms and pressures regarding appearance and norm internalization	Sociocultural Attitudes Toward Appearance Questionnaire-4-Revised [60]
Trait-level ability to eat intuitively following their physical hunger and satiety cues	Intuitive Eating Scale-2 [61]
Fat talk engagement and negative body-related conversations	Fat Talk Questionnaire [62]
Extent to which individuals who engage in disordered eating lie about such behaviors	Deliberate Denial of Disordered Eating Behaviors Scale [63]
Gender experiences and identity	
Experiences of unfair treatment	Everyday Discrimination Scale [64]
Degree to which an individual identifies with and has interest in their own race or ethnicity	Multigroup Ethnic Identity Measure [65]
Degree to which people apply weight-based stereotypes to themselves and base their self-evaluations on weight	Modified Weight Bias Internalization Scale [66]
Sexual minority women’s experiences, identity, and support (for sexual minority women only)	
Sexual minority discrimination	Heterosexist Harassment Rejection, and Discrimination Scale [67]
Sexual minority identity and psychosocial functioning	Lesbian, Gay, and Bisexual Identity Scale [68]
Connectedness to the LGBT ^b community	Connectedness to the LGBT Community Scale [69,70]
Feelings about being part of the sexual minority community and the extent to which their status is important to their identity	Sexual Minority Community Identification [71,72]
Women’s identity and support (for heterosexual women only)	
Feminist identity development	Feminist Identity Composite [73-75]
Individuals’ appraisals of perceived social support	Social Support Appraisals Scale [76]
Mood, stress, and other general life experiences	
Depressive symptoms	Center for Epidemiological Studies Depression 10-item Scale [77]
Anxiety symptoms	Generalized Anxiety Disorder 7-item Scale [78]
Perceived life stress	Perceived Stress Scale [79]
Self-harm and suicidal ideation, plans, and attempts within the past 12 months	Self-Harm Questions-Adapted [80,81]
Ability to bounce back or recover from stress	Brief Resilience Scale [82]
Engagement in comparison behaviors	Iowa-Netherlands Comparison Orientation Measure [83]
Early traumatic experiences and adverse life events	Childhood Traumatic Events Scale [84]
Past 30-day alcohol use—any alcohol use in the last 30 days	N/A
Alcohol use (for women who drank in the last 30 days)	
Quantity, volume, and frequency during a typical week	Daily Drinking Questionnaire [85]

Construct and description	Measure name
Motives for drinking	Drinking Motives Questionnaire [86]
Alcohol consequences experienced by young adult drinkers	Brief Young Adult Alcohol Consequences Questionnaire [87]
Eating-related behaviors related to alcohol consumption	Compensatory Eating and Behaviors in Response to Alcohol Consumption Scale [88,89]
General coping with stress (for women who did not drink in the last 30 days)	
Importance of various reasons for not drinking	Reasons for Not Drinking scale [90]
Strategies for effective and ineffective coping	Brief Coping Orientation to Problems Experienced [91,92]
Other health and health behaviors	
Health-related quality of life	Medical Outcomes Study Short Form-36 [93]
Typical marijuana quantity used each day	General Marijuana Use [85,94]
Typical cigarette quantity used each day	General Cigarette Use [85]
Perceived drinking norms for heterosexual and sexual minority women	Alcohol Descriptive Norms [95]
Reasons why people do not engage in as much physical activity as they think they should	Barriers to Being Active Quiz [96]
Expected outcomes of engaging in physical activity	Exercise Motivation Scale [97]

^aN/A: not applicable (no named measure).

^bLGBT: lesbian, gay, bisexual, and transgender.

Table 2. Ecological momentary assessment surveys.

Survey, construct, and description	Measure name and reference
Morning survey	
Sleep—time in bed, time to fall asleep, wake-up time, length of time asleep, and quality of sleep	Items adapted from the Pittsburgh Sleep Quality Index [98]
Alcohol use—quantity, time drinking, type, location, and interactions (for drinking days only)	Items adapted from various sources, including the study by Heron et al [99]
Compensatory behaviors for drinking (for drinking days only)	Items adapted from the Compensatory Eating and Behaviors in Response to Alcohol Consumption Scale [88]
Nondrinking questions (for nondrinking days only)	Items adapted from various sources, including the study by Heron et al [99]
Mood—anticipated mood for the upcoming day: happy, bored, relaxed, sad, excited, content, worried or anxious, angry, frustrated, or energetic	Items adapted from the Positive and Negative Affect Schedule-Expanded Form [100,101]
Anticipated upcoming stressful and pleasant experiences	Items developed for this study
Before-bed survey	
Mood—overall mood for the day: happy, bored, relaxed, sad, excited, content, worried or anxious, angry, frustrated, or energetic	Items adapted from the Positive and Negative Affect Schedule-Expanded Form [100,101]
Social comparison—nature of social comparison, number of times, type of comparison, person compared with, and feelings about comparison (for social comparison days only)	Items were adapted from sources including the study by Arigo et al [102]
General social experiences—types of people interacted with, pleasantness of interactions, and importance of interactions and topics (for non-social comparison days only)	Items developed for this study
Tobacco—tobacco use	Items developed for this study
Physical activity—length, type, amount, thoughts about activity, estimated amount of physical activity, and exercise the following day	Items adapted from multiple sources [102,103]
Post-binge eating survey	
Date, time, and location of binge eating	Items developed for this study
Eating speed, fullness, with others, and emotions after eating	Items adapted from the Eating Disorder Diagnostic Scale [104] and the DSM-5 ^a binge eating description [105]
Purging and intended purging	Items adapted from the Eating Disorder Diagnostic Scale [104]
Mood before, during, and after binge eating	Items developed for this study based on pilot study participant feedback
Perceived factors contributing to the binge eating episode	Items developed for this study based on pilot study participant feedback
Prompted survey	
Location activity—location of survey notification and activity upon notification	Items developed for this study
Mood—current mood ratings for happy, bored, relaxed, sad, excited, content, worried or anxious, angry, frustrated, or energetic	Affect words selected from the Positive and Negative Affect Schedule-Expanded Form [100,101]
Body image—satisfaction with physical appearance, body shape, weight, physical attractiveness, and looks	Items from the Body Image States Scale [106] adapted from a 9-point scale to a 7-point scale [103]
Self-objectification—thinking about appearance to other people	Item adapted from the study by Holland et al [107]
General stress—stressful or unpleasant experiences	Items adapted from the Daily Inventory of Stressful Events [108] and used in the study by Heron et al [109]
Identity stress—discrimination based on aspect of identity and aspect of identity responsible for discrimination	Items adapted from the study by Panza et al [110]

Survey, construct, and description	Measure name and reference
Disordered eating behaviors—overeating, loss of control of eating, guilt after eating, emotional eating, concerns about eating with others, restriction, avoiding foods	Items from the Eating Disorder Examination Questionnaire 6.0 [111], items adapted using language from the DSM-5 [105], and items from the Emotional Eater Questionnaire [112]
Intuitive eating behaviors—eating for physical hunger, trusting body to eat, body-food choice congruence, permission to eat desirable foods	Items adapted from the Intuitive Eating Scale [61,113]
Physical activity—intensity level of physical activity, type of exercise, length of exercise, and typical amount of exercise	Items adapted from the International Physical Activity Questionnaires [114]
Media use—social media use, type, and activity	Items developed for this study
Appearance-related pressures and source of pressures	Items adapted from the Sociocultural Attitudes Toward Appearance Questionnaire-4 [115]
Negative body-related talk and source of talk	Items adapted from the Fat Talk Questionnaire [62]
Social interactions—type, pleasantness, and importance of people and topic of interaction	Items adapted from the studies by Bernstein et al [116] and Zhaoyan et al [117]

^aDSM-5: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition.

Table 3. End-of-study survey.

Construct and description	Measure name
Accessibility of EMA ^a surveys—questions measuring the burden and accessibility of the EMA surveys	Items developed for this study
Experiences with post-binge eating surveys—measured participants' accurate identification of binge eating episodes and feedback on the survey	Items developed for this study
Identity inclusivity feedback—assessed whether survey items were inclusive of participant identities and captured their experiences	Items developed for this study
Desire to improve health—measured desire and interest in improving physical and mental health	Items developed for this study
Help seeking—assessed resources in social circle that could help the participant address health problems (assessed physical health and mental health problems separately)	Adapted from the General Help-Seeking Questionnaire [118]
Willingness to use mobile health technology—assessed willingness to use mobile health technology to improve physical and mental health or implement health behavior changes	Adapted from the study by Cramer et al [119]
COVID-19 situation—current and previous state of shelter-in-place orders, social distancing practices, and experience of illness with COVID-19	Items developed for this study
COVID-19 health impacts—impact of the COVID-19 pandemic on physical and mental health symptoms as well as change in health behaviors since the start of the pandemic	Items developed for this study [120]
COVID-19 general impacts—changes in responsibilities, finances, and work or school roles since the start of the COVID-19 pandemic	Items selected from the Epidemic-Pandemic Impacts Inventory [121]
COVID-19 minority impacts—impact of the COVID-19 pandemic on minority experiences	Open-ended questions developed for this study

^aEMA: ecological momentary assessment.

Data Analysis Plan

EMA data are inherently hierarchical, with surveys (*level 1*) nested within a person (*level 2*). As such, we will use multilevel modeling to account for similarities between responses for each person. Before hypothesis testing, the data will be examined for normality, outliers, and nonlinear relationships between continuous variables. Model comparison procedures will be used to determine the fixed versus random effects for each model.

Multilevel modeling is robust to missing data (eg, skipped EMA surveys). For variables skipped within a completed assessment, expectation maximization imputation will be used to replace missing values, an approach demonstrated to minimize bias [122]. Participants with complete data will be compared with those with missing data to identify potential attrition biases, and significant predictors will be included as covariates in all analyses.

To test our main hypotheses, level-1 associations between relevant predictors (negative emotions, social factors, and health behaviors) and outcomes (binge eating and loss of control of

eating) will be examined using a series of multilevel models: isolated models with each predictor modeled separately first and then a final model examining all predictors simultaneously to explore unique effects (hypotheses 1a-b). Sexual identity will then be included as a level-2 predictor, and cross-level interactions between sexual orientation (*level 2*) and general factors (*level 1*) will indicate whether momentary associations are different across groups (hypothesis 1c). Aim 2 analyses will use only sexual minority women's data. Level-1 associations between sexual minority stress and eating outcomes will be examined using the same approach as in aim 1: a series of isolated then simultaneous multilevel models (hypothesis 2a). To assess hypothesis 2b, a multilevel structural equation model will be used to examine if negative affect mediates these relationships (1-1-1 model) [123] using Monte Carlo CIs to assess the significance of indirect effects [124]. For aims 1 and 2, we will conduct the aforementioned analyses for momentary associations (random surveys) and event-lagged associations (event and random surveys), allowing for the examination of time-lagged associations (eg, negative affect *before* the event). Aim 3 analyses will examine race- and person-level body- and eating-related factors (BMI, body dissatisfaction, disordered eating, and thin norm internalization), and sexual minority-specific factors (lifetime discrimination, positive sexual minority identity, and community identification and connection) will be included as level-2 predictors in a series of models. Cross-level interactions will indicate whether person-level factors moderate level-1 associations between general factors and binge eating (moderation of aim 1) or between minority-specific factors and binge eating (moderation of aim 2). Three-way interactions among level-1, level-2 sexual orientation, and other level-2 moderators will indicate whether the differential associations detected in the second component of aim 1 (ie, whether associations are stronger or weaker across sexual identity) are moderated by other person-level factors.

Results

Recruitment began in February 2021 and is expected to continue until fall 2022. We have currently consented 129 sexual minority women and 146 heterosexual women into the study. Data cleaning and analysis will begin after data collection is complete.

Discussion

Overview

The goal of the HER Life Project is to examine general and sexual minority-specific predictors of disordered eating, in particular binge eating, among sexual minority and heterosexual young women in their daily lives. This study uses an EMA approach to assess participants at fixed times (morning and before bed), at semirandom times throughout the day, and following self-reported binge eating episodes, thus providing rich information regarding the affective, social, behavioral, and minority-specific experiences of these women. Although there is a growing body of literature considering real-world associations among affective processes (eg, negative affect, stress, and body dissatisfaction), social processes (eg, interactions and social comparisons), and binge eating in general

samples of young women where sexual identity is not known or reported, it remains unclear whether and how these processes operate for sexual minority women and if there are minority-specific predictors (eg, sexual minority stressors) for these women as well. This study includes samples of both sexual minority and heterosexual young women in an effort to explore potential similarities and differences in binge eating experiences and predictors. We expect that there will be many similar affective, social, and health behavior predictors of binge eating across sexual minority and heterosexual women but that there may also be some differences, particularly with regard to body dissatisfaction and unique sexual minority stressors. The findings from this study can help improve our understanding of potential processes that may contribute to health disparities between sexual minority and heterosexual young women's disordered eating behaviors and can be used to inform culturally tailored interventions for binge eating for young sexual minority women. In the following sections, we discuss some of the questions we considered and the challenges we have experienced when designing and carrying out the HER Life Project.

Methodological Challenges

Defining Sexual Identity and Gender Identity

One of the ongoing challenges we experience in our work with sexual minority women is identifying the optimal way to define "sexual minority" and operationalize our definition during the screening process. We initially planned to recruit participants who identified as lesbian women. However, based on some of our previous work [99,125] and in consultation with Community Marketing & Insights, the market research firm we worked with that specializes in recruiting LGBTQ+ adults for web-based research, we learned that young women are increasingly choosing to use other labels when describing their sexual identity (eg, queer or pansexual), use multiple labels (eg, lesbian and queer or gay and queer), or are resisting labels completely [126,127]. Therefore, from the start of this study, we decided to instead focus on recruiting based primarily on sexual attraction and, in some cases, relying also on behavior (ie, gender of the most recent romantic partner). We used a single attraction question that asked people to describe who they were attracted to, with response options including "only or mostly attracted to women," "only or mostly attracted to men," "equally attracted to men and women," or "attracted to people regardless of their gender identity" ("other" and "prefer not to answer" responses were also available). Potential participants were eligible for the heterosexual group if they selected "only or mostly attracted to men." Potential participants were eligible for the sexual minority group if they selected "only or mostly attracted to women" or if they selected "equally attracted" or "attracted regardless of gender" and also endorsed that their current or most recent relationship was with a woman. An ongoing challenge in research on sexual minority individuals is to balance our desire to be inclusive and respectful of the many ways in which identity (including multiple identities) can be defined against our desire to maximize study validity. A sample that is too heterogeneous in terms of identity creates challenges in applying previous literature to our work and interpreting our findings. Therefore, we settled on these criteria in an effort to be more broadly inclusive of sexual minority

women who have diverse attractions (ie, including those with attractions to multiple genders or regardless of gender) but also recognize that, because of the focus in this study on body image and eating behaviors, in the sexual minority women group, we wanted to enroll people who either were attracted to women or were currently or had been recently romantically involved with a woman.

In addition to the challenges in describing sexual identity, we also considered how we defined gender. The focus of this study was on women, and we decided to restrict the sample to people who reported that they were assigned female at birth. We also inquired about potential participants' gender identity, and our initial plan was to only include individuals who identified as "women." However, again in consultation with Community Marketing & Insights, we learned that they are increasingly seeing people identifying with more than one gender label, particularly nonbinary or gender nonconforming. Thus, for this study, we included participants who selected "woman" only or selected both "woman" and either "gender queer/nonconforming" or "nonbinary." Overall, when considering our inclusion criteria for sexual minority and heterosexual women, we tried to balance recruiting relatively homogeneous samples that were distinct from each other based on sexual attraction while also being inclusive of the diverse ways in which young adults describe and understand their identities and attractions [128-130]. These issues of how to define and describe sexual minority individuals will likely be an ongoing challenge for researchers and will continue to evolve as language and cultural understanding change over time.

Remote Data Collection Considerations

Given the large number of young women with diverse sexual identities that we are recruiting for this study, we planned for national recruitment across the United States, which requires that all data collection occur entirely remotely. From the start of the study, we implemented several strategies to enhance study engagement and compliance and also added new protocols throughout based on our experiences. First, to better describe the study to potential participants, we developed several training videos that described the study very generally and others that provided more specific details regarding the study procedures and explained study surveys and materials (eg, defining binge eating). These materials are distributed via email before and during the consent process and are embedded within Qualtrics (Qualtrics International Inc) surveys, allowing us to record whether and for how long the participants review the videos. Participants also have access to several of these relevant videos on the survey app on their phones and can review them throughout the study as needed. On the basis of our pilot work, these videos were well received, and pilot participants described them as useful in orienting them to the intensity of the study procedures so they had a better idea of what to expect when participating.

Second, as part of the screening process, participants shared with us whether they had an Android or Apple phone that they would be willing to use for the study. As a way to confirm potential participants' access to an appropriate smartphone, as part of the screening process (before consent), we ask

participants to download the app to their phones. Although we cannot objectively confirm that they do so, we provide step-by-step downloading instructions as part of a Qualtrics survey and, at the end of the survey, participants indicate that they completed the installation. If participants are unable to install the app on their own phone or do not have a phone that the app is compatible with, we have phones that can be mailed to participants to use for the duration of the study. However, to date, all potential participants have had their own phones to use for the study. This is not surprising given that, in 2021, it was estimated that 96% of adults aged 18 to 29 years owned a smartphone [131]. Together, this suggests that future studies of young adults can expect that most if not all participants will have smartphones that are appropriate for a survey-based EMA study such as this one.

A third procedural consideration that has evolved over the course of the study is how we handle moving people through the study procedure remotely. As described in the Methods section, we developed a reminder system from the start of the study, which includes sending up to 2 reminders, typically several days apart, to participants who are not progressing through the study. Although most participants require very few or no reminders, for those who do receive all the planned reminders from our study team, before either withdrawing them from the study (if they previously consented) or assuming that they are not interested in enrolling (if before consent), the PI sends a final email inquiring about their interest in continuing. Although the PI email does not always result in re-engagement in the study, there have been many instances where it has helped bring a participant back to the study. Overall, this general approach of reminders is helping participants move efficiently through the study process while our study team primarily interacts with participants via email.

The onset of the COVID-19 pandemic and the associated safety protocols and guidelines limited face-to-face interactions in research settings. Given that our study onboarding and data collection procedures were designed to be remote, we experienced a rather seamless transition to conducting our research in the postpandemic era with little to no interruption. Our experiences, coupled with the likelihood of most young adults having a smartphone appropriate for survey-based EMA studies, suggest that entirely remote onboarding and data collection procedures are feasible for national recruitment and in other situations where bringing participants into the laboratory is not always feasible or possible.

Future Research Directions and Study Implications

The HER Life Project primarily focuses on exploring potential similarities and differences in predictors of binge eating between sexual minority and heterosexual young women given the documented health disparities in obesity rates and binge eating between these groups. However, as described in this protocol, part of aim 3 is to explore potential disparities that Black women may experience and consider how racial and sexual minority identities may intersect for Black sexual minority women. As described previously, there is very limited previous research on disordered eating among people who identify as both racial and sexual minority women [46] and, thus, this is a critically

understudied area. Recent reports have highlighted the need for more research using an intersectional perspective and reiterated the importance of such work for providing a more complete understanding of minority health more broadly. Although this study will provide some insight on these topics, it is only a first step, and future work to evaluate the role of other identities (eg, other racial or ethnic group identities, location [urban or rural], and socioeconomic status) and their interaction with sexual orientation is needed to advance the health and well-being of all sexual minority individuals [132]. Continuing to expand and incorporate an intersectional framework will enrich both research and intervention, with ultimate benefits to education, health care, and public policy.

Finally, we view the HER Life Project as 1 step in a large research program aimed at developing culturally sensitive interventions to address binge eating and obesity in sexual minority women. Data from this study can help inform future work in 2 ways. First, as we are including both sexual minority and heterosexual young women, this study will allow us to determine whether there are differential associations between

the hypothesized predictors and binge eating or if sexual minority-specific factors (eg, experiencing sexual minority stressors and discrimination) are associated with binge eating. These findings could help in identifying intervention content for disordered eating or body image interventions that require tailoring for sexual minority women. For example, it is plausible that intervention components teaching strategies for coping with sexual minority stress (instead of engaging in binge eating) are needed. The HER Life Project is also uniquely designed to inform the development of ecological momentary interventions (EMIs). Binge eating EMIs that use mobile apps or SMS text messaging can deliver treatment in women's everyday lives and at specific times when most in need (ie, when likely to engage in binge eating) [133,134]. To develop EMIs, EMA data regarding within-person associations *with the target population* are first needed [135-137], and this study provides such data. In summary, the HER Life Project will provide foundational information needed to develop culturally tailored treatments and mobile technology-based interventions for sexual minority women with the ultimate goal of reducing binge eating and obesity disparities.

Acknowledgments

The research reported in this publication was supported by the National Institute on Minority Health and Health Disparities of the National Institutes of Health under award R01MD012598 to KEH. Author ALB was also supported by an award from the National Institute on Alcohol Abuse and Alcoholism (K01AA023849). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Data Availability

Data will be available from the principal investigator upon reasonable request.

Authors' Contributions

KEH, ALB, and RJL conceptualized the research questions and designed the study and data collection procedures. CAD, CMS, LVB, and AM are responsible for data collection and management. All authors contributed to drafting the manuscript and editing.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Peer-review report from the Psychosocial Risk and Disease Prevention (PRDP) Study, Section - Risk, Prevention and Health Behavior Integrated Review, Group - Center for Scientific Review (National Institutes of Health, USA).

[\[PDF File \(Adobe PDF File\), 179 KB-Multimedia Appendix 1\]](#)

References

1. NHLBI Obesity Education Initiative Expert Panel on the Identification, Evaluation, and Treatment of Obesity in Adults (US). Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. Report No: 98-4083. Bethesda, MD, USA: National Heart, Lung, and Blood Institute; 1998.
2. National Academies of Sciences, Engineering, and Medicine, Division of Behavioral and Social Sciences and Education, Committee on Population, Committee on Understanding the Well-Being of Sexual and Gender Diverse Populations. In: White J, Sepúlveda MJ, Patterson CJ, editors. Understanding the Well-Being of LGBTQI+ Populations. Washington, DC, USA: National Academies Press (US); 2020.
3. Institute of Medicine (US) Committee on Lesbian, Gay, Bisexual, and Transgender Health Issues and Research Gaps and Opportunities. The Health of Lesbian, Gay, Bisexual, and Transgender People: Building a Foundation for Better Understanding. Washington, DC, USA: National Academies Press (US); 2011.
4. Carpenter C. Sexual orientation and body weight: evidence from multiple surveys. *Gend Issues* 2003 Jun;21(3):60-74. [doi: [10.1007/s12147-003-0006-z](https://doi.org/10.1007/s12147-003-0006-z)]

5. Conron KJ, Mimiaga MJ, Landers SJ. A population-based study of sexual orientation identity and gender differences in adult health. *Am J Public Health* 2010 Oct;100(10):1953-1960. [doi: [10.2105/AJPH.2009.174169](https://doi.org/10.2105/AJPH.2009.174169)] [Medline: [20516373](https://pubmed.ncbi.nlm.nih.gov/20516373/)]
6. Struble CB, Lindley LL, Montgomery K, Hardin J, Burcin M. Overweight and obesity in lesbian and bisexual college women. *J Am Coll Health* 2010;59(1):51-56. [doi: [10.1080/07448481.2010.483703](https://doi.org/10.1080/07448481.2010.483703)] [Medline: [20670929](https://pubmed.ncbi.nlm.nih.gov/20670929/)]
7. Austin SB, Ziyadeh NJ, Corliss HL, Rosario M, Wypij D, Haines J, et al. Sexual orientation disparities in purging and binge eating from early to late adolescence. *J Adolesc Health* 2009 Sep;45(3):238-245 [FREE Full text] [doi: [10.1016/j.jadohealth.2009.02.001](https://doi.org/10.1016/j.jadohealth.2009.02.001)] [Medline: [19699419](https://pubmed.ncbi.nlm.nih.gov/19699419/)]
8. Mehl MR, Conner TS. *Handbook of Research Methods for Studying Daily Life*. New York, NY, USA: The Guilford Press; 2013.
9. Smyth JM, Heron KE. Health psychology. In: Mehl MR, Conner TS, editors. *Handbook of Research Methods for Studying Daily Life*. New York, NY, USA: The Guilford Press; 2013:569-584.
10. Smyth J, Wonderlich S, Crosby R, Miltenberger R, Mitchell J, Rorty M. The use of ecological momentary assessment approaches in eating disorder research. *Int J Eat Disord* 2001 Jul;30(1):83-95. [doi: [10.1002/eat.1057](https://doi.org/10.1002/eat.1057)] [Medline: [11439412](https://pubmed.ncbi.nlm.nih.gov/11439412/)]
11. Smyth JM, Wonderlich SA, Heron KE, Sliwinski MJ, Crosby RD, Mitchell JE, et al. Daily and momentary mood and stress are associated with binge eating and vomiting in bulimia nervosa patients in the natural environment. *J Consult Clin Psychol* 2007 Aug;75(4):629-638. [doi: [10.1037/0022-006X.75.4.629](https://doi.org/10.1037/0022-006X.75.4.629)] [Medline: [17663616](https://pubmed.ncbi.nlm.nih.gov/17663616/)]
12. Haedt-Matt AA, Keel PK. Revisiting the affect regulation model of binge eating: a meta-analysis of studies using ecological momentary assessment. *Psychol Bull* 2011 Jul;137(4):660-681 [FREE Full text] [doi: [10.1037/a0023660](https://doi.org/10.1037/a0023660)] [Medline: [21574678](https://pubmed.ncbi.nlm.nih.gov/21574678/)]
13. Mason TB, Lewis RJ, Heron KE. Daily discrimination and binge eating among lesbians: a pilot study. *Psychol Sex* 2017 Mar 04;8(1-2):96-103. [doi: [10.1080/19419899.2017.1296484](https://doi.org/10.1080/19419899.2017.1296484)]
14. Hawkins RC, Clement PF. Binge eating: measurement problems and a conceptual model. In: Hawkins RC, Fremouw WJ, Clement PF, editors. *The Binge-Purge Syndrome: Diagnosis, Treatment, and Research*. New York, NY, USA: Springer Publishing Company; 1984:229-251.
15. Srivastava P, Michael ML, Manasse SM, Juarascio AS. Do momentary changes in body dissatisfaction predict binge eating episodes? An ecological momentary assessment study. *Eat Weight Disord* 2021 Feb;26(1):395-400. [doi: [10.1007/s40519-020-00849-z](https://doi.org/10.1007/s40519-020-00849-z)] [Medline: [31989487](https://pubmed.ncbi.nlm.nih.gov/31989487/)]
16. Jones MD, Crowther JH, Ciesla JA. A naturalistic study of fat talk and its behavioral and affective consequences. *Body Image* 2014 Sep;11(4):337-345. [doi: [10.1016/j.bodyim.2014.05.007](https://doi.org/10.1016/j.bodyim.2014.05.007)] [Medline: [24976570](https://pubmed.ncbi.nlm.nih.gov/24976570/)]
17. Leahey TM, Crowther JH, Ciesla JA. An ecological momentary assessment of the effects of weight and shape social comparisons on women with eating pathology, high body dissatisfaction, and low body dissatisfaction. *Behav Ther* 2011 Jun;42(2):197-210 [FREE Full text] [doi: [10.1016/j.beth.2010.07.003](https://doi.org/10.1016/j.beth.2010.07.003)] [Medline: [21496506](https://pubmed.ncbi.nlm.nih.gov/21496506/)]
18. Mills J, Fuller-Tyszkiewicz M, Holmes M. State body dissatisfaction and social interactions: an experience sampling study. *Psychol Women Q* 2014 Feb 03;38(4):551-562. [doi: [10.1177/0361684314521139](https://doi.org/10.1177/0361684314521139)]
19. Morrison MA, Morrison TG, Sager C. Does body satisfaction differ between gay men and lesbian women and heterosexual men and women? A meta-analytic review. *Body Image* 2004 May;1(2):127-138. [doi: [10.1016/j.bodyim.2004.01.002](https://doi.org/10.1016/j.bodyim.2004.01.002)] [Medline: [18089146](https://pubmed.ncbi.nlm.nih.gov/18089146/)]
20. Levine MP, Smolak L. *The Prevention of Eating Problems and Eating Disorders: Theories, Research, and Applications*. Mahwah, NJ, USA: Lawrence Erlbaum Associates; 2006.
21. Stice E. Risk and maintenance factors for eating pathology: a meta-analytic review. *Psychol Bull* 2002 Sep;128(5):825-848. [doi: [10.1037/0033-2909.128.5.825](https://doi.org/10.1037/0033-2909.128.5.825)] [Medline: [12206196](https://pubmed.ncbi.nlm.nih.gov/12206196/)]
22. Thompson JK, Stice E. Thin-ideal internalization: mounting evidence for a new risk factor for body-image disturbance and eating pathology. *Curr Dir Psychol Sci* 2001 Oct 1;10(5):181-183. [doi: [10.1111/1467-8721.00144](https://doi.org/10.1111/1467-8721.00144)]
23. VanKim NA, Porta CM, Eisenberg ME, Neumark-Sztainer D, Laska MN. Lesbian, gay and bisexual college student perspectives on disparities in weight-related behaviours and body image: a qualitative analysis. *J Clin Nurs* 2016 Dec;25(23-24):3676-3686 [FREE Full text] [doi: [10.1111/jocn.13106](https://doi.org/10.1111/jocn.13106)] [Medline: [27878900](https://pubmed.ncbi.nlm.nih.gov/27878900/)]
24. Fingerhut AW, Peplau LA, Ghavami N. A dual-identity framework for understanding lesbian experience. *Psychol Women Q* 2005;29(2):129-139. [doi: [10.1111/j.1471-6402.2005.00175.x](https://doi.org/10.1111/j.1471-6402.2005.00175.x)]
25. Gadalla T, Piran N. Co-occurrence of eating disorders and alcohol use disorders in women: a meta analysis. *Arch Womens Ment Health* 2007;10(4):133-140. [doi: [10.1007/s00737-007-0184-x](https://doi.org/10.1007/s00737-007-0184-x)] [Medline: [17533558](https://pubmed.ncbi.nlm.nih.gov/17533558/)]
26. Mason TB, Lewis RJ. Minority stress, depression, relationship quality, and alcohol use: associations with overweight and obesity among partnered young adult lesbians. *LGBT Health* 2015 Dec;2(4):333-340 [FREE Full text] [doi: [10.1089/lgbt.2014.0053](https://doi.org/10.1089/lgbt.2014.0053)] [Medline: [26788774](https://pubmed.ncbi.nlm.nih.gov/26788774/)]
27. Vancampfort D, Vanderlinden J, Stubbs B, Soundy A, Pieters G, De Hert M, et al. Physical activity correlates in persons with binge eating disorder: a systematic review. *Eur Eat Disord Rev* 2014 Jan;22(1):1-8. [doi: [10.1002/erv.2255](https://doi.org/10.1002/erv.2255)] [Medline: [24014460](https://pubmed.ncbi.nlm.nih.gov/24014460/)]
28. Crandall KJ, Eisenman PA, Ransdell L, Reel J. Exploring binge eating and physical activity among community-dwelling women. *Int J Psychol Behav Sci* 2011 Nov 31;1(1):1-8 [FREE Full text] [doi: [10.5923/j.ijpbs.20110101.01](https://doi.org/10.5923/j.ijpbs.20110101.01)]

29. Yancey AK, Cochran SD, Corliss HL, Mays VM. Correlates of overweight and obesity among lesbian and bisexual women. *Prev Med* 2003 Jun;36(6):676-683 [FREE Full text] [doi: [10.1016/s0091-7435\(03\)00020-3](https://doi.org/10.1016/s0091-7435(03)00020-3)] [Medline: [12744910](https://pubmed.ncbi.nlm.nih.gov/12744910/)]
30. Drabble L, Midanik LT, Trocki K. Reports of alcohol consumption and alcohol-related problems among homosexual, bisexual and heterosexual respondents: results from the 2000 National Alcohol Survey. *J Stud Alcohol* 2005 Jan;66(1):111-120. [doi: [10.15288/jsa.2005.66.111](https://doi.org/10.15288/jsa.2005.66.111)] [Medline: [15830911](https://pubmed.ncbi.nlm.nih.gov/15830911/)]
31. McCabe SE, Hughes TL, Bostwick WB, West BT, Boyd CJ. Sexual orientation, substance use behaviors and substance dependence in the United States. *Addiction* 2009 Aug;104(8):1333-1345 [FREE Full text] [doi: [10.1111/j.1360-0443.2009.02596.x](https://doi.org/10.1111/j.1360-0443.2009.02596.x)] [Medline: [19438839](https://pubmed.ncbi.nlm.nih.gov/19438839/)]
32. Wilsnack SC, Hughes TL, Johnson TP, Bostwick WB, Szalacha LA, Benson P, et al. Drinking and drinking-related problems among heterosexual and sexual minority women. *J Stud Alcohol Drugs* 2008 Jan;69(1):129-139. [doi: [10.15288/jsad.2008.69.129](https://doi.org/10.15288/jsad.2008.69.129)] [Medline: [18080073](https://pubmed.ncbi.nlm.nih.gov/18080073/)]
33. Calzo JP, Roberts AL, Corliss HL, Blood EA, Kroshus E, Austin SB. Physical activity disparities in heterosexual and sexual minority youth ages 12-22 years old: roles of childhood gender nonconformity and athletic self-esteem. *Ann Behav Med* 2014 Feb;47(1):17-27 [FREE Full text] [doi: [10.1007/s12160-013-9570-y](https://doi.org/10.1007/s12160-013-9570-y)] [Medline: [24347406](https://pubmed.ncbi.nlm.nih.gov/24347406/)]
34. Laska MN, VanKim NA, Erickson DJ, Lust K, Eisenberg ME, Rosser BR. Disparities in weight and weight behaviors by sexual orientation in college students. *Am J Public Health* 2015 Jan;105(1):111-121. [doi: [10.2105/AJPH.2014.302094](https://doi.org/10.2105/AJPH.2014.302094)] [Medline: [25393177](https://pubmed.ncbi.nlm.nih.gov/25393177/)]
35. Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence. *Psychol Bull* 2003 Sep;129(5):674-697 [FREE Full text] [doi: [10.1037/0033-2909.129.5.674](https://doi.org/10.1037/0033-2909.129.5.674)] [Medline: [12956539](https://pubmed.ncbi.nlm.nih.gov/12956539/)]
36. Hatzenbuehler ML. How does sexual minority stigma "get under the skin"? A psychological mediation framework. *Psychol Bull* 2009 Sep;135(5):707-730 [FREE Full text] [doi: [10.1037/a0016441](https://doi.org/10.1037/a0016441)] [Medline: [19702379](https://pubmed.ncbi.nlm.nih.gov/19702379/)]
37. Ancheta AJ, Caceres BA, Zollweg SS, Heron KE, Veldhuis CB, VanKim NA, et al. Examining the associations of sexual minority stressors and past-year depression with overeating and binge eating in a diverse community sample of sexual minority women. *Eat Behav* 2021 Dec;43:101547. [doi: [10.1016/j.eatbeh.2021.101547](https://doi.org/10.1016/j.eatbeh.2021.101547)] [Medline: [34412003](https://pubmed.ncbi.nlm.nih.gov/34412003/)]
38. Watson LB, Grotewiel M, Farrell M, Marshik J, Schneider M. Experiences of sexual objectification, minority stress, and disordered eating among sexual minority women. *Psychol Women Q* 2015 Mar 02;39(4):458-470. [doi: [10.1177/0361684315575024](https://doi.org/10.1177/0361684315575024)]
39. Mason TB, Lewis RJ. Minority stress and binge eating among lesbian and bisexual women. *J Homosex* 2015;62(7):971-992. [doi: [10.1080/00918369.2015.1008285](https://doi.org/10.1080/00918369.2015.1008285)] [Medline: [25603175](https://pubmed.ncbi.nlm.nih.gov/25603175/)]
40. Panza E, Fehling KB, Pantalone DW, Dodson S, Selby EA. Multiply marginalized: linking minority stress due to sexual orientation, gender, and weight to dysregulated eating among sexual minority women of higher body weight. *Psychol Sex Orientat Gend Divers* 2021 Dec;8(4):420-428. [doi: [10.1037/sgd0000431](https://doi.org/10.1037/sgd0000431)] [Medline: [34926715](https://pubmed.ncbi.nlm.nih.gov/34926715/)]
41. Heffernan K. Eating disorders and weight concern among lesbians. *Int J Eat Disord* 1996 Mar;19(2):127-138. [doi: [10.1002/\(SICI\)1098-108X\(199603\)19:2<127::AID-EAT3>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1098-108X(199603)19:2<127::AID-EAT3>3.0.CO;2-P)] [Medline: [8932551](https://pubmed.ncbi.nlm.nih.gov/8932551/)]
42. Striegel-Moore RH, Tucker N, Hsu J. Body image dissatisfaction and disordered eating in lesbian college students. *Int J Eat Disord* 1990 Sep;9(5):493-500. [doi: [10.1002/1098-108X\(199009\)9:5<493::AID-EAT2260090504>3.0.CO;2-C](https://doi.org/10.1002/1098-108X(199009)9:5<493::AID-EAT2260090504>3.0.CO;2-C)]
43. Fitzgibbon ML, Spring B, Avellone ME, Blackman LR, Pingitore R, Stolley MR. Correlates of binge eating in Hispanic, Black, and White women. *Int J Eat Disord* 1998 Jul;24(1):43-52. [doi: [10.1002/\(sici\)1098-108x\(199807\)24:1<43::aid-eat4>3.0.co;2-0](https://doi.org/10.1002/(sici)1098-108x(199807)24:1<43::aid-eat4>3.0.co;2-0)] [Medline: [9589310](https://pubmed.ncbi.nlm.nih.gov/9589310/)]
44. Howard LM, Heron KE, MacIntyre RI, Myers TA, Everhart RS. Is use of social networking sites associated with young women's body dissatisfaction and disordered eating? A look at Black-White racial differences. *Body Image* 2017 Dec;23:109-113. [doi: [10.1016/j.bodyim.2017.08.008](https://doi.org/10.1016/j.bodyim.2017.08.008)] [Medline: [28965051](https://pubmed.ncbi.nlm.nih.gov/28965051/)]
45. Pike KM, Dohm FA, Striegel-Moore RH, Wilfley DE, Fairburn CG. A comparison of black and white women with binge eating disorder. *Am J Psychiatry* 2001 Sep;158(9):1455-1460. [doi: [10.1176/appi.ajp.158.9.1455](https://doi.org/10.1176/appi.ajp.158.9.1455)] [Medline: [11532731](https://pubmed.ncbi.nlm.nih.gov/11532731/)]
46. Feldman MB, Meyer IH. Eating disorders in diverse lesbian, gay, and bisexual populations. *Int J Eat Disord* 2007 Apr;40(3):218-226 [FREE Full text] [doi: [10.1002/eat.20360](https://doi.org/10.1002/eat.20360)] [Medline: [17262818](https://pubmed.ncbi.nlm.nih.gov/17262818/)]
47. Herek GM, Garnets LD. Sexual orientation and mental health. *Annu Rev Clin Psychol* 2007;3:353-375. [doi: [10.1146/annurev.clinpsy.3.022806.091510](https://doi.org/10.1146/annurev.clinpsy.3.022806.091510)] [Medline: [17716060](https://pubmed.ncbi.nlm.nih.gov/17716060/)]
48. Fingerhut A, Peplau LA, Gable SL. Identity, minority stress and psychological well-being among gay men and lesbians. *Psychol Sex* 2010 Jun 03;1(2):101-114. [doi: [10.1080/19419899.2010.484592](https://doi.org/10.1080/19419899.2010.484592)]
49. Doty ND, Willoughby BL, Lindahl KM, Malik NM. Sexuality related social support among lesbian, gay, and bisexual youth. *J Youth Adolesc* 2010 Oct;39(10):1134-1147. [doi: [10.1007/s10964-010-9566-x](https://doi.org/10.1007/s10964-010-9566-x)] [Medline: [20593304](https://pubmed.ncbi.nlm.nih.gov/20593304/)]
50. O'Hara RE, Armeli S, Scott DM, Covault J, Tennen H. Perceived racial discrimination and negative-mood-related drinking among African American college students. *J Stud Alcohol Drugs* 2015 Mar;76(2):229-236 [FREE Full text] [doi: [10.15288/jsad.2015.76.229](https://doi.org/10.15288/jsad.2015.76.229)] [Medline: [25785798](https://pubmed.ncbi.nlm.nih.gov/25785798/)]
51. Wood SM, Schott W, Marshal MP, Akers AY. Disparities in body mass index trajectories from adolescence to early adulthood for sexual minority women. *J Adolesc Health* 2017 Dec;61(6):722-728 [FREE Full text] [doi: [10.1016/j.jadohealth.2017.06.008](https://doi.org/10.1016/j.jadohealth.2017.06.008)] [Medline: [28935384](https://pubmed.ncbi.nlm.nih.gov/28935384/)]

52. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007 May;39(2):175-191. [doi: [10.3758/bf03193146](https://doi.org/10.3758/bf03193146)] [Medline: [17695343](https://pubmed.ncbi.nlm.nih.gov/17695343/)]
53. West SG, Ryu E, Kwok OM, Cham H. Multilevel modeling: current and future applications in personality research. *J Pers* 2011 Feb;79(1):2-50. [doi: [10.1111/j.1467-6494.2010.00681.x](https://doi.org/10.1111/j.1467-6494.2010.00681.x)] [Medline: [21223263](https://pubmed.ncbi.nlm.nih.gov/21223263/)]
54. Heron KE, Braitman AL, Lewis RJ, Shappie AT, Hitson PT. Measuring sexual minority stressors in lesbian women's daily lives: initial scale development. *Psychol Sex Orientat Gend Divers* 2018 Sep;5(3):387-395 [FREE Full text] [doi: [10.1037/sgd0000287](https://doi.org/10.1037/sgd0000287)] [Medline: [30221178](https://pubmed.ncbi.nlm.nih.gov/30221178/)]
55. Heron KE, Scott SB, Sliwinski MJ, Smyth JM. Eating behaviors and negative affect in college women's everyday lives. *Int J Eat Disord* 2014 Dec;47(8):853-859 [FREE Full text] [doi: [10.1002/eat.22292](https://doi.org/10.1002/eat.22292)] [Medline: [24797029](https://pubmed.ncbi.nlm.nih.gov/24797029/)]
56. Fritz MS, Mackinnon DP. Required sample size to detect the mediated effect. *Psychol Sci* 2007 Mar;18(3):233-239 [FREE Full text] [doi: [10.1111/j.1467-9280.2007.01882.x](https://doi.org/10.1111/j.1467-9280.2007.01882.x)] [Medline: [17444920](https://pubmed.ncbi.nlm.nih.gov/17444920/)]
57. Forbush KT, Wildes JE, Pollack LO, Dunbar D, Luo J, Patterson K, et al. Development and validation of the Eating Pathology Symptoms Inventory (EPSI). *Psychol Assess* 2013 Sep;25(3):859-878. [doi: [10.1037/a0032639](https://doi.org/10.1037/a0032639)] [Medline: [23815116](https://pubmed.ncbi.nlm.nih.gov/23815116/)]
58. Mond JM, Hay PJ, Rodgers B, Owen C. Eating Disorder Examination Questionnaire (EDE-Q): norms for young adult women. *Behav Res Ther* 2006 Jan;44(1):53-62. [doi: [10.1016/j.brat.2004.12.003](https://doi.org/10.1016/j.brat.2004.12.003)] [Medline: [16301014](https://pubmed.ncbi.nlm.nih.gov/16301014/)]
59. Cooper PJ, Taylor MJ, Cooper Z, Fairbum CG. The development and validation of the body shape questionnaire. *Int J Eat Disord* 1987 Jul;6(4):485-494. [doi: [10.1002/1098-108X\(198707\)6:4<485::AID-EAT2260060405>3.0.CO;2-O](https://doi.org/10.1002/1098-108X(198707)6:4<485::AID-EAT2260060405>3.0.CO;2-O)]
60. Schaefer LM, Harriger JA, Heinberg LJ, Soderberg T, Kevin Thompson J. Development and validation of the sociocultural attitudes towards appearance questionnaire-4-revised (SATAQ-4R). *Int J Eat Disord* 2017 Feb;50(2):104-117. [doi: [10.1002/eat.22590](https://doi.org/10.1002/eat.22590)] [Medline: [27539814](https://pubmed.ncbi.nlm.nih.gov/27539814/)]
61. Tylka TL. Development and psychometric evaluation of a measure of intuitive eating. *J Couns Psychol* 2006 Apr;53(2):226-240. [doi: [10.1037/0022-0167.53.2.226](https://doi.org/10.1037/0022-0167.53.2.226)]
62. Royal S, Macdonald DE, Dionne MM. Development and validation of the Fat Talk Questionnaire. *Body Image* 2013 Jan;10(1):62-69. [doi: [10.1016/j.bodyim.2012.10.003](https://doi.org/10.1016/j.bodyim.2012.10.003)] [Medline: [23201392](https://pubmed.ncbi.nlm.nih.gov/23201392/)]
63. Howard LM, Heron KE, Cramer RJ. The deliberate denial of disordered eating behaviors scale: development and initial validation in young women with subclinical disordered eating. *J Psychopathol Behav Assess* 2020 Jun 18;42(4):774-786 [FREE Full text] [doi: [10.1007/s10862-020-09819-2](https://doi.org/10.1007/s10862-020-09819-2)]
64. Williams DR, Yu Y, Jackson JS, Anderson NB. Racial differences in physical and mental health: socio-economic status, stress and discrimination. *J Health Psychol* 1997 Jul;2(3):335-351. [doi: [10.1177/135910539700200305](https://doi.org/10.1177/135910539700200305)] [Medline: [22013026](https://pubmed.ncbi.nlm.nih.gov/22013026/)]
65. Phinney JS. The Multigroup Ethnic Identity Measure: a new scale for use with diverse groups. *J Adoles Res* 1992 Apr 1;7(2):156-176. [doi: [10.1177/074355489272003](https://doi.org/10.1177/074355489272003)]
66. Pearl RL, Puhl RM. Measuring internalized weight attitudes across body weight categories: validation of the modified weight bias internalization scale. *Body Image* 2014 Jan;11(1):89-92. [doi: [10.1016/j.bodyim.2013.09.005](https://doi.org/10.1016/j.bodyim.2013.09.005)] [Medline: [24100004](https://pubmed.ncbi.nlm.nih.gov/24100004/)]
67. Szymanski DM. Does internalized heterosexism moderate the link between heterosexist events and lesbians' psychological distress? *Sex Roles* 2006 Feb;54(3-4):227-234. [doi: [10.1007/s11199-006-9340-4](https://doi.org/10.1007/s11199-006-9340-4)]
68. Mohr JJ, Kendra MS. Revision and extension of a multidimensional measure of sexual minority identity: the Lesbian, Gay, and Bisexual Identity Scale. *J Couns Psychol* 2011 Apr;58(2):234-245. [doi: [10.1037/a0022858](https://doi.org/10.1037/a0022858)] [Medline: [21319899](https://pubmed.ncbi.nlm.nih.gov/21319899/)]
69. Frost DM, Meyer IH. Measuring community connectedness among diverse sexual minority populations. *J Sex Res* 2012;49(1):36-49 [FREE Full text] [doi: [10.1080/00224499.2011.565427](https://doi.org/10.1080/00224499.2011.565427)] [Medline: [21512945](https://pubmed.ncbi.nlm.nih.gov/21512945/)]
70. Zimmerman L, Darnell DA, Rhew IC, Lee CM, Kaysen D. Resilience in community: a social ecological development model for young adult sexual minority women. *Am J Community Psychol* 2015 Mar;55(1-2):179-190 [FREE Full text] [doi: [10.1007/s10464-015-9702-6](https://doi.org/10.1007/s10464-015-9702-6)] [Medline: [25572956](https://pubmed.ncbi.nlm.nih.gov/25572956/)]
71. Boyle SC, Omoto AM. Lesbian community oughts and ideals: normative fit, depression, and anxiety among young sexual minority women. *Psychol Women Q* 2014;38(1):33-45. [doi: [10.1177/0361684313484900](https://doi.org/10.1177/0361684313484900)]
72. Luhtanen R, Crocker J. A collective self-esteem scale: self-evaluation of one's social identity. *Pers Soc Psychol Bull* 1992 Jun 1;18(3):302-318. [doi: [10.1177/0146167292183006](https://doi.org/10.1177/0146167292183006)]
73. Fischer AR, Tokar DM, Mergl MM, Good GE, Hill MS, Blum SA. Assessing women's feminist identity development: studies of convergent, discriminant, and structural validity. *Psychol Women Q* 2000 Mar;24(1):15-29. [doi: [10.1111/j.1471-6402.2000.tb01018.x](https://doi.org/10.1111/j.1471-6402.2000.tb01018.x)]
74. Bargad A, Hyde JS. Women's studies: a study of feminist identity development in women. *Psychol Women Q* 1991 Jun 1;15(2):181-201. [doi: [10.1111/j.1471-6402.1991.tb00791.x](https://doi.org/10.1111/j.1471-6402.1991.tb00791.x)]
75. Rickard KM. The relationship of self-monitored dating behaviors to level of feminist identity on the feminist identity scale. *Sex Roles* 1989 Feb;20(3-4):213-226. [doi: [10.1007/BF00287993](https://doi.org/10.1007/BF00287993)]
76. Vaux A, Phillips J, Holly L, Thomson B, Williams D, Stewart D. The social support appraisals (SS-A) scale: studies of reliability and validity. *Am J Commun Psychol* 1986 Apr;14(2):195-218. [doi: [10.1007/BF00911821](https://doi.org/10.1007/BF00911821)]

77. Andresen EM, Malmgren JA, Carter WB, Patrick DL. Screening for depression in well older adults: evaluation of a short form of the CES-D (Center for Epidemiologic Studies Depression Scale). *Am J Prev Med* 1994;10(2):77-84. [Medline: [8037935](#)]
78. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006 May 22;166(10):1092-1097 [FREE Full text] [doi: [10.1001/archinte.166.10.1092](#)] [Medline: [16717171](#)]
79. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav* 1983 Dec;24(4):385-396. [Medline: [6668417](#)]
80. Nock MK, Holmberg EB, Photos VI, Michel BD. Self-injurious thoughts and behaviors interview: development, reliability, and validity in an adolescent sample. *Psychol Assess* 2007 Sep;19(3):309-317. [doi: [10.1037/1040-3590.19.3.309](#)] [Medline: [17845122](#)]
81. Miller GK, Piscopo KD, Batts K, Han B, Colpe L, Forman-Hoffman VL, et al. Measurement of suicidal thoughts, behaviors, and related health outcomes in the United States: comparison of NSDUH estimates with other data sources. Substance Abuse and Mental Health Services Administration. 2015 Jul. URL: <https://www.samhsa.gov/data/sites/default/files/NSDUH-DR-N20Suicide-2015/NSDUH-DR-N20Suicide-2015.pdf> [accessed 2020-10-15]
82. Smith BW, Dalen J, Wiggins K, Tooley E, Christopher P, Bernard J. The brief resilience scale: assessing the ability to bounce back. *Int J Behav Med* 2008;15(3):194-200. [doi: [10.1080/10705500802222972](#)] [Medline: [18696313](#)]
83. Gibbons FX, Buunk BP. Individual differences in social comparison: development of a scale of social comparison orientation. *J Pers Soc Psychol* 1999 Jan;76(1):129-142. [doi: [10.1037//0022-3514.76.1.129](#)] [Medline: [9972558](#)]
84. Pennebaker JW, Susman JR. Disclosure of traumas and psychosomatic processes. *Soc Sci Med* 1988;26(3):327-332. [doi: [10.1016/0277-9536\(88\)90397-8](#)] [Medline: [3279521](#)]
85. Collins RL, Parks GA, Marlatt GA. Social determinants of alcohol consumption: the effects of social interaction and model status on the self-administration of alcohol. *J Consult Clin Psychol* 1985 Apr;53(2):189-200. [doi: [10.1037//0022-006x.53.2.189](#)] [Medline: [3998247](#)]
86. Cooper ML. Motivations for alcohol use among adolescents: development and validation of a four-factor model. *Psychol Assess* 1994 Jun;6(2):117-128 [FREE Full text] [doi: [10.1037/1040-3590.6.2.117](#)]
87. Kahler CW, Strong DR, Read JP. Toward efficient and comprehensive measurement of the alcohol problems continuum in college students: the brief young adult alcohol consequences questionnaire. *Alcohol Clin Exp Res* 2005 Jul;29(7):1180-1189. [doi: [10.1097/01.alc.0000171940.95813.a5](#)] [Medline: [16046873](#)]
88. Rahal CJ, Bryant JB, Darkes J, Menzel JE, Thompson JK. Development and validation of the Compensatory Eating and Behaviors in Response to Alcohol Consumption Scale (CEBRACS). *Eat Behav* 2012 Apr;13(2):83-87. [doi: [10.1016/j.eatbeh.2011.11.001](#)] [Medline: [22365787](#)]
89. Choquette EM, Dedrick R, Thompson JK, Rancourt D. Reexamination of the psychometric properties of the Compensatory Eating and Behaviors in Response to Alcohol Consumption Scale (CEBRACS) and exploration of alternative scoring. *Eat Behav* 2020 Aug;38:101410. [doi: [10.1016/j.eatbeh.2020.101410](#)] [Medline: [32736310](#)]
90. Johnson TJ, Cohen EA. College students' reasons for not drinking and not playing drinking games. *Subst Use Misuse* 2004 Jun;39(7):1137-1160. [doi: [10.1081/ja-120038033](#)] [Medline: [15387207](#)]
91. Carver CS. You want to measure coping but your protocol's too long: consider the brief COPE. *Int J Behav Med* 1997;4(1):92-100. [doi: [10.1207/s15327558ijbm0401_6](#)] [Medline: [16250744](#)]
92. Carver CS, Scheier MF, Weintraub JK. Assessing coping strategies: a theoretically based approach. *J Pers Soc Psychol* 1989 Feb;56(2):267-283. [doi: [10.1037//0022-3514.56.2.267](#)] [Medline: [2926629](#)]
93. Ware Jr JJ, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992 Jun;30(6):473-483. [Medline: [1593914](#)]
94. Collins RL, Vincent PC, Yu J, Liu L, Epstein LH. A behavioral economic approach to assessing demand for marijuana. *Exp Clin Psychopharmacol* 2014 Jun;22(3):211-221 [FREE Full text] [doi: [10.1037/a0035318](#)] [Medline: [24467370](#)]
95. Ehlike SJ, Stamates AL, Kelley ML, Braitman AL. Bisexual women's reports of descriptive drinking norms for heterosexual, bisexual, and lesbian women. *Psychol Sex Orientat Gend Divers* 2019 Jun;6(2):256-263 [FREE Full text] [doi: [10.1037/sgd0000312](#)] [Medline: [31106227](#)]
96. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Division of Nutrition and Physical Activity. Barriers to being active quiz. In: Brown DR, Heath G, Martin SL, editors. *Promoting Physical Activity: A Guide for Community Action*. Champaign, IL, USA: Human Kinetics; 1999:100-101.
97. Steinhardt MA, Dishman RK. Reliability and validity of expected outcomes and barriers for habitual physical activity. *J Occup Med* 1989 Jun;31(6):536-546. [doi: [10.1097/00043764-198906000-00011](#)] [Medline: [2786559](#)]
98. Buysse DJ, Reynolds 3rd CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 1989 May;28(2):193-213. [doi: [10.1016/0165-1781\(89\)90047-4](#)] [Medline: [2748771](#)]
99. Heron KE, Lewis RJ, Shappie AT, Dawson CA, Amerson R, Braitman AL, et al. Rationale and design of a remote Web-based daily diary study examining sexual minority stress, relationship factors, and alcohol use in same-sex female couples across

- the United States: study protocol of project relate. *JMIR Res Protoc* 2019 Feb 04;8(2):e11718 [FREE Full text] [doi: [10.2196/11718](https://doi.org/10.2196/11718)] [Medline: [30714946](https://pubmed.ncbi.nlm.nih.gov/30714946/)]
100. Watson D, Clark LA. The PANAS-X: Manual for the Positive and Negative Affect Schedule - Expanded Form. The University of Iowa. 1994. URL: <https://www2.psychology.uiowa.edu/faculty/clark/panas-x.pdf> [accessed 2020-09-30]
 101. Posner J, Russell JA, Peterson BS. The circumplex model of affect: an integrative approach to affective neuroscience, cognitive development, and psychopathology. *Dev Psychopathol* 2005;17(3):715-734 [FREE Full text] [doi: [10.1017/S0954579405050340](https://doi.org/10.1017/S0954579405050340)] [Medline: [16262989](https://pubmed.ncbi.nlm.nih.gov/16262989/)]
 102. Arigo D, Brown MM, Pasko K, Ainsworth MC, Travers L, Gupta A, et al. Rationale and design of the women's health and daily experiences project: protocol for an ecological momentary assessment study to identify real-time predictors of midlife women's physical activity. *JMIR Res Protoc* 2020 Oct 15;9(10):e19044 [FREE Full text] [doi: [10.2196/19044](https://doi.org/10.2196/19044)] [Medline: [33055065](https://pubmed.ncbi.nlm.nih.gov/33055065/)]
 103. MacIntyre RI, Heron KE, Braitman AL, Arigo D. An ecological momentary assessment of self-improvement and self-evaluation body comparisons: associations with college women's body dissatisfaction and exercise. *Body Image* 2020 Jun;33:264-277 [FREE Full text] [doi: [10.1016/j.bodyim.2020.04.002](https://doi.org/10.1016/j.bodyim.2020.04.002)] [Medline: [32473545](https://pubmed.ncbi.nlm.nih.gov/32473545/)]
 104. Stice E, Telch CF, Rizvi SL. Development and validation of the eating disorder diagnostic scale: a brief self-report measure of anorexia, bulimia, and binge-eating disorder. *Psychol Assess* 2000 Jun;12(2):123-131. [doi: [10.1037//1040-3590.12.2.123](https://doi.org/10.1037//1040-3590.12.2.123)] [Medline: [10887758](https://pubmed.ncbi.nlm.nih.gov/10887758/)]
 105. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders (DSM-5). 5th edition. Washington, DC, USA: American Psychiatric Association; 2013.
 106. Cash TF, Fleming EC, Alindogan J, Steadman L, Whitehead A. Beyond body image as a trait: the development and validation of the Body Image States Scale. *Eat Disord* 2002;10(2):103-113. [doi: [10.1080/10640260290081678](https://doi.org/10.1080/10640260290081678)] [Medline: [16864251](https://pubmed.ncbi.nlm.nih.gov/16864251/)]
 107. Holland E, Koval P, Stratemeyer M, Thomson F, Haslam N. Sexual objectification in women's daily lives: a smartphone ecological momentary assessment study. *Br J Soc Psychol* 2017 Jun;56(2):314-333. [doi: [10.1111/bjso.12152](https://doi.org/10.1111/bjso.12152)] [Medline: [27484394](https://pubmed.ncbi.nlm.nih.gov/27484394/)]
 108. Almeida DM, Wethington E, Kessler RC. The daily inventory of stressful events: an interview-based approach for measuring daily stressors. *Assessment* 2002 Mar;9(1):41-55. [doi: [10.1177/1073191102091006](https://doi.org/10.1177/1073191102091006)] [Medline: [11911234](https://pubmed.ncbi.nlm.nih.gov/11911234/)]
 109. Heron KE, Scott SB, Mogle JA, Howard LM, Everhart RS. Ambulatory assessment of everyday stressors: a two-study experiment evaluating the effect of question format on self-reported daily and momentary stressors. *J Technol Behav Sci* 2022 May 07;7(3):368-380 [FREE Full text] [doi: [10.1007/s41347-022-00259-1](https://doi.org/10.1007/s41347-022-00259-1)]
 110. Panza E, Olson K, Goldstein CM, Selby EA, Lillis J. Characterizing lifetime and daily experiences of weight stigma among sexual minority women with overweight and obesity: a descriptive study. *Int J Environ Res Public Health* 2020 Jul 07;17(13):4892 [FREE Full text] [doi: [10.3390/ijerph17134892](https://doi.org/10.3390/ijerph17134892)] [Medline: [32645883](https://pubmed.ncbi.nlm.nih.gov/32645883/)]
 111. Fairburn CG, Beglin S. Eating disorder examination questionnaire (EDE-Q6.0). In: Fairburn CG, editor. *Cognitive Behavior Therapy and Eating Disorders*. New York, NY, USA: Guilford Press; 2008:309-313.
 112. Garaulet M, Canteras M, Morales E, López-Guimera G, Sánchez-Carracedo D, Corbalán-Tutau MD. Validation of a questionnaire on emotional eating for use in cases of obesity: the Emotional Eater Questionnaire (EEQ). *Nutr Hosp* 2012;27(2):645-651. [doi: [10.1590/S0212-16112012000200043](https://doi.org/10.1590/S0212-16112012000200043)] [Medline: [22732995](https://pubmed.ncbi.nlm.nih.gov/22732995/)]
 113. Tylka TL, Kroon Van Diest AM. The Intuitive Eating Scale-2: item refinement and psychometric evaluation with college women and men. *J Couns Psychol* 2013 Jan;60(1):137-153. [doi: [10.1037/a0030893](https://doi.org/10.1037/a0030893)] [Medline: [23356469](https://pubmed.ncbi.nlm.nih.gov/23356469/)]
 114. Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 2003 Aug;35(8):1381-1395. [doi: [10.1249/01.MSS.0000078924.61453.FB](https://doi.org/10.1249/01.MSS.0000078924.61453.FB)] [Medline: [12900694](https://pubmed.ncbi.nlm.nih.gov/12900694/)]
 115. Schaefer LM, Burke NL, Thompson JK, Dedrick RF, Heinberg LJ, Calogero RM, et al. Development and validation of the Sociocultural Attitudes Towards Appearance Questionnaire-4 (SATAQ-4). *Psychol Assess* 2015 Mar;27(1):54-67. [doi: [10.1037/a0037917](https://doi.org/10.1037/a0037917)] [Medline: [25285718](https://pubmed.ncbi.nlm.nih.gov/25285718/)]
 116. Bernstein MJ, Zawadzki MJ, Juth V, Benfield JA, Smyth JM. Social interactions in daily life: within-person associations between momentary social experiences and psychological and physical health indicators. *J Soc Pers Relationships* 2017 Feb 16;35(3):372-394 [FREE Full text] [doi: [10.1177/0265407517691366](https://doi.org/10.1177/0265407517691366)]
 117. Zhaoyang R, Sliwinski MJ, Martire LM, Smyth JM. Age differences in adults' daily social interactions: an ecological momentary assessment study. *Psychol Aging* 2018 Jun;33(4):607-618 [FREE Full text] [doi: [10.1037/pag0000242](https://doi.org/10.1037/pag0000242)] [Medline: [29708385](https://pubmed.ncbi.nlm.nih.gov/29708385/)]
 118. Wilson CJ, Deane FP, Ciarrochi JV, Rickwood D. Measuring help seeking intentions: Properties of the General Help Seeking Questionnaire. University of Wollongong. 2005. URL: <https://ro.uow.edu.au/cgi/viewcontent.cgi?referer=&httpsredir=1&article=2580&context=hbspapers> [accessed 2020-10-18]
 119. Cramer RJ, Wright S, Wilsey CN, Kaniuka AR, Bowling J, Crocker T, et al. Alternative sexuality, sexual orientation and mobile technology: findings from the National Coalition for Sexual Freedom technology and health enhancement feasibility study. *Psychol Sex* 2022;13(2):344-359 [FREE Full text] [doi: [10.1080/19419899.2020.1777188](https://doi.org/10.1080/19419899.2020.1777188)]
 120. Whittaker F, Kingston S. Stress, social support, and substance use in the COVID-19 pandemic. *Transl Issues Psychol Sci* (forthcoming) 2022 Apr 14. [doi: [10.1037/tps0000322](https://doi.org/10.1037/tps0000322)]

121. Grasso DJ, Briggs-Gowan MJ, Ford JD, Carter AS. Epidemic – Pandemic Impacts Inventory (EPII). University of Connecticut School of Medicine. 2020. URL: https://www.phenxtoolkit.org/toolkit_content/PDF/Grasso_EPII.pdf [accessed 2020-09-16]
122. Allison PD. Missing data. In: Quantitative Applications in the Social Sciences. Thousand Oaks, CA, USA: Sage Publications; 2001.
123. Preacher KJ, Zyphur MJ, Zhang Z. A general multilevel SEM framework for assessing multilevel mediation. *Psychol Methods* 2010 Sep;15(3):209-233. [doi: [10.1037/a0020141](https://doi.org/10.1037/a0020141)] [Medline: [20822249](https://pubmed.ncbi.nlm.nih.gov/20822249/)]
124. Preacher KJ, Selig JP. Advantages of Monte Carlo confidence intervals for indirect effects. *Commun Methods Meas* 2012 Apr;6(2):77-98. [doi: [10.1080/19312458.2012.679848](https://doi.org/10.1080/19312458.2012.679848)]
125. Lewis RJ, Dawson CA, Shappie AT, Braitman AL, Heron KE. Recruiting cisgender female couples for health disparity-focused daily diary research: challenges, successes, and lessons learned. *Psychol Sex* (forthcoming) 2021 Jun 23:1-21 [FREE Full text] [doi: [10.1080/19419899.2021.1942177](https://doi.org/10.1080/19419899.2021.1942177)]
126. Savin-Williams R. Refusing and resisting sexual identity labels. In: Browning DL, editor. *Adolescent Identities: A Collection of Readings*. Milton Park, UK: Taylor & Francis; 2008:67-91.
127. Galupo MP, Mitchell RC, Davis KS. Sexual minority self-identification: multiple identities and complexity. *Psychol Sex Orientat Gend Divers* 2015 Dec;2(4):355-364 [FREE Full text] [doi: [10.1037/sgd0000131](https://doi.org/10.1037/sgd0000131)]
128. Savin-Williams RC, Joyner K, Rieger G. Prevalence and stability of self-reported sexual orientation identity during young adulthood. *Arch Sex Behav* 2012 Feb;41(1):103-110. [doi: [10.1007/s10508-012-9913-y](https://doi.org/10.1007/s10508-012-9913-y)] [Medline: [22302504](https://pubmed.ncbi.nlm.nih.gov/22302504/)]
129. Talley AE, Stevens JE. Sexual orientation self-concept ambiguity: scale adaptation and validation. *Assessment* 2017 Jul;24(5):632-645 [FREE Full text] [doi: [10.1177/1073191115617016](https://doi.org/10.1177/1073191115617016)] [Medline: [26643117](https://pubmed.ncbi.nlm.nih.gov/26643117/)]
130. Young RM, Meyer IH. The trouble with "MSM" and "WSW": erasure of the sexual-minority person in public health discourse. *Am J Public Health* 2005 Jul;95(7):1144-1149. [doi: [10.2105/AJPH.2004.046714](https://doi.org/10.2105/AJPH.2004.046714)] [Medline: [15961753](https://pubmed.ncbi.nlm.nih.gov/15961753/)]
131. Mobile Fact Sheet. Pew Research Center. 2021 Apr. URL: <https://www.pewresearch.org/internet/fact-sheet/mobile/> [accessed 2022-06-29]
132. Else-Quest NM, Hyde JS. Intersectionality in quantitative psychological research: I. Theoretical and epistemological issues. *Psychol Women Q* 2016 Feb 26;40(2):155-170. [doi: [10.1177/0361684316629797](https://doi.org/10.1177/0361684316629797)]
133. Heron KE, Smyth JM. Ecological momentary interventions: incorporating mobile technology into psychosocial and health behaviour treatments. *Br J Health Psychol* 2010 Feb;15(Pt 1):1-39 [FREE Full text] [doi: [10.1348/135910709X466063](https://doi.org/10.1348/135910709X466063)] [Medline: [19646331](https://pubmed.ncbi.nlm.nih.gov/19646331/)]
134. Juarascio AS, Goldstein SP, Manasse SM, Forman EM, Butryn ML. Perceptions of the feasibility and acceptability of a smartphone application for the treatment of binge eating disorders: qualitative feedback from a user population and clinicians. *Int J Med Inform* 2015 Oct;84(10):808-816 [FREE Full text] [doi: [10.1016/j.ijmedinf.2015.06.004](https://doi.org/10.1016/j.ijmedinf.2015.06.004)] [Medline: [26113461](https://pubmed.ncbi.nlm.nih.gov/26113461/)]
135. Borrelli B, Ritterband LM. Special issue on eHealth and mHealth: challenges and future directions for assessment, treatment, and dissemination. *Health Psychol* 2015 Dec;34S:1205-1208. [doi: [10.1037/hea0000323](https://doi.org/10.1037/hea0000323)] [Medline: [26651461](https://pubmed.ncbi.nlm.nih.gov/26651461/)]
136. Smith KE, Juarascio A. From ecological momentary assessment (EMA) to ecological momentary intervention (EMI): past and future directions for ambulatory assessment and interventions in eating disorders. *Curr Psychiatry Rep* 2019 Jun 04;21(7):53. [doi: [10.1007/s11920-019-1046-8](https://doi.org/10.1007/s11920-019-1046-8)] [Medline: [31161276](https://pubmed.ncbi.nlm.nih.gov/31161276/)]
137. Spruijt-Metz D, Nilsen W. Dynamic models of behavior for just-in-time adaptive interventions. *IEEE Pervasive Comput* 2014 Jul;13(3):13-17. [doi: [10.1109/MPRV.2014.46](https://doi.org/10.1109/MPRV.2014.46)]

Abbreviations

- EMA:** ecological momentary assessment
 - EMI:** ecological momentary intervention
 - HER Life:** Health and Experiences in Real Life
 - ICC:** intraclass correlation coefficient
 - LGBTQ+:** lesbian, gay, bisexual, transgender, and queer
 - PI:** principal investigator
-

Edited by T Leung; This paper was peer reviewed by the Psychosocial Risk and Disease Prevention (PRDP) Study, Section - Risk, Prevention and Health Behavior Integrated Review, Group - Center for Scientific Review (National Institutes of Health, USA). See the Multimedia Appendix for the peer-review report; Submitted 19.07.22; accepted 30.08.22; published 21.10.22.

Please cite as:

Heron KE, Braitman AL, Dawson CA, Sandoval CM, Butler LV, Moulder A, Lewis RJ

Rationale and Design of an Ecological Momentary Assessment Study Examining Predictors of Binge Eating Among Sexual Minority and Heterosexual Young Women: Protocol for the Health and Experiences in Real Life (HER Life) Study

JMIR Res Protoc 2022;11(10):e41199

URL: <https://www.researchprotocols.org/2022/10/e41199>

doi: [10.2196/41199](https://doi.org/10.2196/41199)

PMID:

©Kristin E Heron, Abby L Braitman, Charlotte A Dawson, Cassidy M Sandoval, Lauren V Butler, Alicia Moulder, Robin J Lewis. Originally published in JMIR Research Protocols (<https://www.researchprotocols.org>), 21.10.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Research Protocols, is properly cited. The complete bibliographic information, a link to the original publication on <https://www.researchprotocols.org>, as well as this copyright and license information must be included.